



STUDIA UNIVERSITATIS
BABEŞ-BOLYAI



BIOLOGIA

1/2018

**STUDIA
UNIVERSITATIS BABEŞ-BOLYAI
BIOLOGIA**

**1 / 2018
January – June**

EDITORIAL BOARD

STUDIA UNIVERSITATIS BABEȘ-BOLYAI BIOLOGIA

EDITOR-IN-CHIEF:

Professor **Octavian Popescu**, Member of the Romanian Academy, Babeș-Bolyai University, Cluj-Napoca.

BOARD OF SUBJECT EDITORS:

Associate Professor **Ioan Coroiu**, Babeș-Bolyai University, Cluj-Napoca;
Professor **Nicolae Dragoș**, Babeș-Bolyai University, Cluj-Napoca;
Professor **László Gallé**, Member of the Hungarian Academy, University of Szeged, Hungary;
Professor **Michael Moustakas**, Aristotle University, Thessaloniki, Greece;
Professor **Aharon Oren**, Alexander Silberman Institute of Life Sciences, Jerusalem, Israel;
Professor **Leontin Ștefan Péterfi**, Associate Member of the Romanian Academy, Babeș-Bolyai University, Cluj-Napoca;
Professor **László Rakosy**, Babeș-Bolyai University, Cluj-Napoca;
Senior Researcher **Anca Sima**, Associate Member of the Romanian Academy, Institute of Citology and Cellular Pathology, Bucharest;
Professor **Helga Stan-Lötter**, University of Salzburg, Salzburg, Austria;
Professor **Corneliu Tarba**, Babeș-Bolyai University, Cluj-Napoca.

LIST OF ASSOCIATE REVIEWERS:

Senior Researcher **Adela Halmagyi**, Institute of Biological Research, Cluj-Napoca;
Professor **László Rakosy**, Babeș-Bolyai University, Cluj-Napoca;
Associate Professor **Ioan Coroiu**, Babeș-Bolyai University, Cluj-Napoca;
Lecturer **Rahela Carpa**, Babeș-Bolyai University, Cluj-Napoca;
Lecturer **Alin David**, Babeș-Bolyai University, Cluj-Napoca;
Lecturer **Anca Farkaș**, Babeș-Bolyai University, Cluj-Napoca;
Teaching Assistant **Cristina Mircea**, Babeș-Bolyai University, Cluj-Napoca;
Biologist **Sára Ferentî**, University of Oradea, Faculty of Sciences, Oradea.

EXECUTIVE EDITORS:

Lecturer **Karina Paula Battes**, Babeș-Bolyai University, Cluj-Napoca;
Lecturer **Mirela Cîmpean**, Babeș-Bolyai University, Cluj-Napoca.
Contact: studies.ubb.biologia@gmail.com

YEAR
MONTH
ISSUE

Volume 63 (LXIII) 2018
JUNE
1

PUBLISHED ONLINE: 2018-06-22
PUBLISHED PRINT: 2018-06-29
ISSUE DOI:10.24193/subbbiol.2018.1

STUDIA
UNIVERSITATIS BABEȘ-BOLYAI
BIOLOGIA

1

STUDIA UBB EDITORIAL OFFICE: B.P. Hasdeu no. 51, 400371 Cluj-Napoca, Romania,
Phone + 40 264 405352, www.studia.ubbcluj.ro

SUMAR – CONTENTS – SOMMAIRE – INHALT

REGULAR ARTICLES

- V. BULIMAGA, M. PISOVA, L. ZOSIM, A. TROFIM, Procedures of partial purification for phycobiliproteins from cyanobacteria isolated from soils of Republic of Moldova 5
- N. TOMESCU, L. A. TEODOR, Terrestrial isopods (Isopoda, Crustacea) from the “Danube Delta” Biosphere Reserve 15
- A. MESTECĂNEANU, R. GAVA, Ornithological observations on the Golești Basin (Argeș River, Romania) between February 2013 and January 2014..... 25

E. E. IMARHIAGBE, B. IKHAJIAGBE, Antibiotic susceptibility of bacterial isolates and water quality index of water sourced from closed ground water and open hand dug well in Koko Community, Delta State, Nigeria	47
A. -L. BUTIUC-KEUL, L. JARDA, I. GOIA, I. HOLOBIUC, A. FARKAS, V. CRISTEA, Preliminary data regarding genetic diversity of several endangered and endemic <i>Dianthus</i> species from Romania generated by RAPD markers	59
N. TOMESCU, L. A. TEODOR, <i>Protracheoniscus vasileradui</i> – n. sp. (Crustacea, Isopoda, Crinochaeta) in the Romanian fauna	73
A. DAVID, A. N. STERMIN, E. SEVIANU, Clutch size and egg repeatability in three elusive bird species: Little Bittern (<i>Ixobrychus minutus</i>), Little Crake (<i>Zapornia parva</i>) and Water Rail (<i>Rallus aquaticus</i>) from north-west Romanian populations	81

All authors are responsible for submitting manuscripts in comprehensible US or UK English and ensuring scientific accuracy.

Original pictures on front cover: *Protracheoniscus vasileradui* n. sp. (Crustacea, Isopoda, Crinochaeta) (male) © Nicolae Tomescu and Lucian Alexandru Teodor

Procedures of partial purification for phycobiliproteins from cyanobacteria isolated from soils of Republic of Moldova

Valentina Bulimaga¹, Maria Pisova¹, Liliana Zosim¹ and Alina Trofim¹✉

SUMMARY. Investigation of the new cyanobacterial strains, for use as potential sources of bioactive substances, including phycobiliproteins, encounters some difficulties due to presence of toxins (microcystins) produced by some cyanobacterial strains. Cyanobacteria phycobiliproteins are natural pigments with high potential for application as colorants in food, cosmetics and pharmaceuticals. The objective of the study was the elaboration of a procedure for *Anabaena propinqua* Setchell. et Gardn. phycobiliproteins separation from microcystins and a procedure of partial purification of phycobiliproteins from cyanobacteria *Anabenopsis* sp. The antioxidant capacity of partial purified phycoerythrin from *Anabenopsis* sp. was established.

Keywords: antioxidant capacity, cyanobacteria, microcystins, phycobiliproteins.

Introduction

Cyanobacteria possess a wide spectrum of actual and potential biotechnological applications in diverse fields, such as agriculture, aquaculture, bioremediation, bioenergy and biofuels, nutraceuticals and pharmaceuticals, food industry, cosmetics and biomedical research (Abed *et al.*, 2009; Chu, 2012; Lau *et al.*, 2015; Manirafasha *et al.*, 2016).

Investigation of the new cyanobacterial strains for use as potential sources of bioactive substances encounters some difficulties with the presence of toxins, including microcystins, produced by some cyanobacteria. Most microcystins are hepatotoxins (liver toxins). Hepatotoxins are produced by species of the genera *Microcystis*, *Anabaena*, *Nodularia*, *Oscillatoria*, *Cylindrospermum* (Bulimaga *et al.*,

¹ SRL "Phycobiotecology", Moldova State University, Chișinău, Republic of Moldova, 65A. M. Kogălniceanu Street, MD 2009.

✉ **Corresponding author: Alina Trofim**, Moldova State University, Chișinău, Republic of Moldova, 5A. M. Kogălniceanu Street, MD 2009,
E-mail: alinatrofim@yahoo.com

2014). Moreover, toxins can be eliminated in the nutritive media or can be extracted together with bioactive substances. The methods used for removing of microcystins from drinking water are mainly based on application of activated carbon (Pyo and Moon, 2005; Yan *et al.*, 2006; Drogui *et al.*, 2012).

At the same time, publications regarding the removal of microcystins from phycobiliproteins extracts are in very small number (Ehmann and Guthrie, 2011, 2015).

The goal of the present research was to elaborate the procedures for removal of microcystins from *Anabaena propinqua* phycobiliprotein extracts and partial purification of phycoerythrin from cyanobacteria *Anabaenopsis* sp.

Materials and methods

The strains of investigated cyanobacteria *Anabaena propinqua* Setchell. et Gardn. and *Anabaenopsis* sp. (isolated by A. Trofim) were cultivated and offered by SRL “Algology”, State University of Moldova, under the leadership of the professor V. M. Șalaru. The strains were isolated from the soils of the Cogalnic River Valley meadow, Cimișlia, Republic of Moldova.

Separation of phycobiliproteins extracted from biomass of cyanobacteria *Anabaena propinqua* from toxins was performed by the chromatographic method. The Amberlite XAD-2 (Sigma-Aldrich) column (20 x 0.5 cm) was washed with 10 volumes–of bidistilled water. The aqueous suspension of *Anabaena propinqua* biomass (20 mg/ml) was supposed to freeze-thawed repeated procedure and subsequent maceration of the frozen mixture, using pestle in a mortar for 1 min. The macerate was centrifuged at 6000 rpm, 10 minutes. Extract (42 ml) was placed in the Amberlite column. The unabsorbed fraction was eluted with bidistilled H₂O and the toxin-free phycobilliproteins solution was obtained. Toxins (microcystins) were adsorbed on the Amberlite and could be eluted with alcoholic solutions. To elute the toxins adsorbed on Amberlite, the column was washed with H₂O, 20% C₂H₅OH, then with 96% C₂H₅OH. The identification of the toxin fractions was performed at 240 nm. The peptidic nature of microcystins was established by reaction with 0.35% ninhydrin.

Partial purification of phycobiliproteins extracts from cyanobacteria *Anabaenopsis* sp. by (NH₄)₂SO₄ precipitation. Fractionation of phycobiliproteins was carried out by (NH₄)₂SO₄ precipitation (Pandey *et al.*, 2011; Chakdar and Pabbi, 2012) with our modification. To 70 ml 9.92 g (NH₄)₂SO₄ were added to 25% saturation and after storage 1 hour at 4°C the suspension was centrifuged at 10000 rpm, 10 minutes. A pink precipitate containing phycoerythrin was obtained. The precipitate was dissolved in 15 ml of water and after centrifugation the supernatant - (phycoerythrin 1) was collected and the insoluble residue (pink violet precipitate) poorly soluble in water has been removed. Then (NH₄)₂SO₄ was added to the supernatant to

60% saturation and, after 1 hour of storage at 4°C, the sample was centrifuged at 10000 rpm, 10 min. The precipitate was dissolved in H₂O and the insoluble residue was removed by centrifugation. The pink violet supernatant was collected (phycoerythrin 3), and the pink residue was dissolved in 1.5 ml H₂O and centrifuged. In the obtained supernatant phycoerythrin 2 was isolated. The preparations of phycoerythrin were further subjected to dialysis for 24 hours against 100 times volume of MilliQ water containing 3 mM sodium azide.

Antioxidant activity assessment of phycoerythrin preparations by ABTS radical cation scavenging assay (Re *et al.*, 1999). Antioxidant activity of phycoerythrin 1 isolated by 25% (NH₄)₂SO₄ fractionation, as well as the fractions obtained by 25-60% (NH₄)₂SO₄ precipitation - phycoerythrin 2 and 3 was determined by the reaction with the cation ABTS⁺ (2,2-azinobis-3-ethylbenzothiazoline-6-sulfonic acid). ABTS⁺ was generated by oxidation of ABTS with potassium persulfate. 7 mM ABTS solution and potassium persulfate (2.45 mM) were dissolved in deionized water. The reaction mixture was preserved at the room temperature for 12-16 hours in the dark before using. ABTS⁺ from the stock solution was diluted with ethanol to absorbance at 734 nm of 0.700 ± 0.020. Then 1 ml of diluted ABTS⁺ solution was mixed with 0.1 or 0.3 ml of the test sample (1.0 mg/ml) and after 6 min the absorbance was measured at 734 nm.

The % inhibition was calculated according to the equation:

$$\% \text{ Inhibition} = \left[\frac{Abs_{t0} - Abs_{t6}}{Abs_{t0}} \right] \times 100 \%$$

where Abs_{t0 min} is the extinction value of the ABTS⁺ and Abs_{t6min} solution is the extinction value of the ABTS⁺ solution after 6 min of incubation with the samples. All determinations were performed in 3 replicates.

Results and discussion

Removing of microcystins from *Anabaena propinqua* phycobiliproteins extract. In the present study the *Anabaena propinqua* phycobiliproteins extract has been analyzed. Freezing and thawing method and subsequent maceration of the frozen mixture, using pestle in a mortar, were selected as efficient way to obtain aqueous extract of phycobiliproteins from *Anabaena propinqua*. The extraction of phycobiliproteins was accompanied by the presence of toxins (microcystins).

Separation of toxins (microcystins) from phycobiliproteins was performed by the chromatographic method on Amberlite XAD-2 (Fig.1). Toxins (microcystins) were adsorbed on the Amberlite and could be eluted with alcoholic solutions. To elute the toxins adsorbed on Amberlite, the column was washed with H₂O, 20% C₂H₅OH, then with 96% C₂H₅OH. The identification of the toxin fractions was performed at 240 nm. The peptide nature of microcystins was established by reaction with 0.35% ninhydrin.




Figure 1. Scheme of microcystins removing from *Anabaena propinqua* phycobiliproteins extract.

Partial purification of *Anabaenopsis* sp. phycobiliproteins by two steps $(\text{NH}_4)_2\text{SO}_4$ precipitation. As a result of the partial purification of phycobiliproteins aqueous extract by two steps fractionation with $(\text{NH}_4)_2\text{SO}_4$ (Table 1, Fig. 2) three preparations of phycoerythrin were obtained: phycoerythrin 1 from precipitate isolated after 25% $(\text{NH}_4)_2\text{SO}_4$ fractionation, phycoerythrin 3 from precipitate obtained by 25-60% $(\text{NH}_4)_2\text{SO}_4$ fractionation and phycoerythrin 2 from residue resulted after solubilization of the precipitate obtained by 25-60% $(\text{NH}_4)_2\text{SO}_4$ fractionation.

Table 1.

Partial purification of phycobiliproteins extracts of cyanobacteria *Anabaenopsis* sp by $(\text{NH}_4)_2\text{SO}_4$ precipitation

Phycoerythrin fractions	A565	A620	A650	A280	Phycoerythrin purity (A565/A280)
Phycobiliproteins extract	0.400	0.212	0.120	0.400	1.0
Phycoerythrin 1	0.881	0.184	0.122	0.569	1.54
Phycoerythrin 2	1.045	0.220	0.142	0.528	1.98
Phycoerythrin 3	1.210	0.826	0.097	0.481	2.5

Scheme of partial purification of *Anabaenopsis* sp. phycobiliproteins by two steps $(\text{NH}_4)_2\text{SO}_4$ precipitation is presented in Fig. 2.




Figure 2. Scheme of partial purification of *Anabaenopsis* sp. phycobiliproteins by two steps $(\text{NH}_4)_2\text{SO}_4$ precipitation

It has been established that the two consecutive steps of phycobiliproteins purification by $(\text{NH}_4)_2\text{SO}_4$ precipitation was efficient for partial purification of phycoerythrin from *Anabaenopsis* sp. aqueous extract (Table 1, Fig. 3a, b).

UV-VIS absorbance of phycoerythrin preparations obtained from *Anabaenopsis* sp. extract at the first and second step of fractionation by $(\text{NH}_4)_2\text{SO}_4$ revealed that the maximum content of phycoerythrin with the highest purity ($A_{565}/A_{280}=2.5$) was detected in the phycoerythrin fraction obtained by 25-60% $(\text{NH}_4)_2\text{SO}_4$ precipitation (Fig. 3b). The preparation contains phycocyanin besides phycoerythrin. The purity of phycoerythrin 1 is the lowest ($A_{565}/A_{280}=1.54$) in comparison with the other phycoerythrin preparations. A high absorbance at 280 nm was observed, that can be connected with presence of ballast proteins in this solution (Fig. 3a).




Figure 3. UV-VIS absorbance of phycoerythrin preparations obtained from *Anabaenopsis sp* extract at the 1-st and second step of purification by $(\text{NH}_4)_2\text{SO}_4$: a) phycoerythrin 1 (PE1) and phycoerythrin 2(PE2); b) phycoerythrin 3 (PE3), containing both phycoerythrin and phycocyanin (PC)

Antioxidant capacity of phycoerythrin preparations obtained from cyanobacteria Anabaenopsis sp. assessed by ABTS⁺ method.

Consumption of natural antioxidants, such as phycobiliproteins, that can possibly scavenge free radicals has often been referred as an effective therapeutic option to alleviate free radicals induced cellular damage. Oxidative stress plays a key role in onset and progression of pathophysiological manifestation of many diseases, including cancer. Intra-cellular oxidative stress takes place under conditions of production of excessive ROS that cannot be mitigated by antioxidant defense system.

Partial purified phycoerythrin preparations obtained from cyanobacteria *Anabaenopsis sp.* were tested for determination of its antioxidant capacity (Table 2). The obtained results allow us to conclude that the phycoerythrin 3 fraction constituted from phycoerythrin and phycocyanin, obtained from cyanobacterium *Anabaenopsis sp.* after partial purification by 25-60% $(\text{NH}_4)_2\text{SO}_4$ precipitation, possesses a maximum antioxidant capacity (100%). In case of phycoerythrin 1 and phycoerythrin 2 lower values of antioxidant capacity (24.9 and 27.48%, respectively) are recorded.

Table 2.

Antioxidant capacity of phycoerythrin preparations obtained from cyanobacteria *Anabaena* sp. determined by ABTS⁺ method

Sample	A734	A734(0) - 734(exp)	% inhibition	
			0.1 ml	0.3ml
Phycoerythrin fraction after the 1-st step of fractionation (25% (NH ₄) ₂ SO ₄)				
1. ABTS	0.699±0.04			
2. Phycoerythrin 1	0.641±0.03	0.058	8.30	24.90
Phycoerythrin fractions after the 2second step of fractionation (25-60%(NH ₄) ₂ SO ₄)				
3. Phycoerythrin 2	0.635±0.03	0.064	9.16	27.48
4. Phycoerythrin 3	0.464±0.02	0.235	33.62	100

From previous research it was established that cyanobacterium *Anabaena propinqua* contains 7.22–8.87% of phycobiliproteins from biomass, at the cultivation on the Drew media supplemented with NH₄NO₃. So, the phycoerythrin content was prevailing (4.29 – 5.10% of biomass) in comparison with phycocyanin and allophycocyanin content. The content of phycocyanin and allophycocyanin varies between 0.53 to 2.3% and 0.67 to 2.09%, respectively (Bulimaga *et al.*, 2014). The high content of phycobiliproteins (up to 8.87%) in cyanobacterium *Anabaena propinqua* biomass makes it a source of perspective for obtaining natural colorants.

The extraction of phycobiliproteins can be carried out using phosphate buffers or distilled water. However, the water extraction is more preferable having the advantage of phycobiliproteins obtaining with a higher yield compared to extraction with buffers (Khatoona *et al.*, 2018). The method for rapid phycobiliproteins extraction from cyanobacteria *Synechococcus* CCMP 833 is also known (Viskari and Colyer, 2003). The disadvantage of this method is the necessity in dialysis of phycobiliproteins extract for removing of detergent CHAPS, used in high concentration (3%) for culture cells disruption.

The use of *Anabaena propinqua* as a source of phycobiliproteins is limited due to the presence of microcystins that are extracted together with phycobiliproteins. The research carried out in the present paper has shown that AmberliteXAD-2 (hydrophobic copolymer of styrene-divinylbenzene resin) can be used as efficient adsorbent for microcystins.

The toxin-free phycobilliproteins fraction was not adsorbed on Amberlite column and could be eluted by H₂O. For the separation of microcystins from phycobiliproteins, the authors Ehmman and Guthrie have used other resins, such as

Amberlite™ XAD 16HP, Amberlite™ FPX66, Diaion™ PS-DVB or Sepabeads™ SP70 (Ehmman and Guthrie, 2011, 2015).

As a result of purification of *Anabaenopsis* sp. phycobiliproteins by two steps fractionation with $(\text{NH}_4)_2\text{SO}_4$, three fractions of phycoerythrin: PE-1, PE-2, and PE-3 were obtained. According to the purity values (A_{620}/A_{280}) of phycoerythrin fractions, the purest fraction is PE-3(2.5) followed by PE-2(1.98) and PE-1(1.54) (Table 1).

Although the PE-3 fraction had a higher purity and could be used as a food and cosmetic pigment, it also contains phycocyanin, besides phycoerythrin (Fig. 3b). Further purification of the PE-3 fraction could be performed by chromatographic methods for use in immunodiagnosics or drug preparations. The other two phycoerythrin fractions contained a higher amount of ballast protein (Fig. 3a).

The scheme proposed in this study can be used to fractionate and obtain partially purified phycoerythrin fraction not only at *Anabaenopsis* sp. and *Anabaena propinqua*, but also to other cyanobacteria.

The analysis of the antioxidant activity of the obtained phycoerythrin fractions revealed the maximum antioxidant capacity of phycoerythrin 3, containing some quantity of phycocyanin. This fact is probably due to their synergistic action. The results are in accordance with the research results of various authors who related the high antioxidant capacity of C-phycocyanin from cyanobacteria *Spirulina platensis* (Bulimaga *et al.*, 2012) and *Synechococcus* sp. (Sonani *et al.*, 2017), as well as C-phycoerythrin from *Phormidium* sp. and *Halomicronema* sp. (Madamwar *et al.*, 2015).

Conclusions

Separation of *Anabaena propinqua* phycobiliproteins from toxins (microcystins) by the chromatographic method on Amberlite was performed. The unabsorbed fraction was eluted with distilled H_2O and the toxins-free phycobilliproteins solution was obtained. The procedure of phycobiliproteins isolation from microcystins was proposed. It has been established that the two consecutive steps of purification of phycoerythrin by $(\text{NH}_4)_2\text{SO}_4$ precipitation were efficient for partial purification of phycobiliproteins from *Anabaenopsis* sp. aqueous extract. The maximum content of phycoerythrin with the highest purity ($A_{565}/A_{280}=2.5$) was detected in the phycoerythrin fraction obtained by 25-60% $(\text{NH}_4)_2\text{SO}_4$ precipitation. The antioxidant capacity of phycoerythrin preparations has been established.

REFERENCES

- Abed, R. M., Dobretsov, S., Sudesh, K. (2009) Applications of cyanobacteria in biotechnology, *J. Appl. Microbiol.* **106** (1):1-12
- Bulimaga, V., Djur, S., Pisov, M., Rudi, L. Rudic, V. (2012) Antioxidant capacity of phycocyanin preparations obtained from Ge-enriched *Spirulina* biomass, *Studia Universitatis Moldaviae, Științe reale și ale naturii*, **1** (51): 9-13 [in Romanian]
- Bulimaga, V., Șalaru V. M., Zosim, L., Pisov, M., Trofim, A. (2014) Biochemical composition of the blue-green algae *Anabaena propinqua* at cultivation on the Drew nutritive media with and without nitrogen sources, *2nd International Conference on microbial technology*. October 9-10, Chisinau Moldova, pp.106
- Bulimaga, V., Șalaru, V., Zosim, L., Pisov, M., Trofim, A. (2014) Blue-green algae (Cyanophyta) - sources of bioactive secondary metabolites, *Studia Universitatis Moldaviae, Științe reale și ale naturii*, **1** (71): 96-107 [in Romanian]
- Chakdar, H., Pabbi, S. (2012) Extraction and purification of phycoerythrin from *Anabaena variabilis* (CCC421), *Phykos*, **42** (1): 25 – 31
- Chu, W. -L. (2012) Biotechnological applications of microalgae, *Je JSME.*, **6**(Suppl1): S24-S37
- Drogui, P., Daghrrir, R., Simard, M.C., Sauvageau, C., Blais, J. F. (2012) Removal of microcystin-LR from spiked water using either activated carbon or anthracite as filter material, *Environ Technol.*, **33**(4-6):381-91
- Ehmann, A., Guthrie, J. (2011) Methods for removal of microcystins and isolation of phycocyanin from *Cyanobacteria*. Patent. WO2011011174, Publication date 27.01.2011
- Ehmann, A., Guthrie, J. (2015) Methods for removal of microcystins and isolation of phycocyanin from cyanobacteria, Patent US 9131724 B2, Publication date 15.09.2015
- Khatona , H., Leong, L. K. , Rahman N. A., Mian S., Begum, H., Banerjee S. Endut, A. (2018) Effects of different light source and media on growth and production of phycobiliprotein from freshwater cyanobacteria, *Bioresource Technology*, **249**: 652–658
- Lau, N. -S., Matsui, M., Abdullah, A. A. -A. (2015) Cyanobacteria: Photoautotrophic Microbial Factories for the Sustainable Synthesis of Industrial Products, BioMed Research International, Article ID 754934, <http://dx.doi.org/10.1155/2015/754934>
- Madamwar, D., Patel, D. K., Desai, S. N., Upadhyay, K. K., Devkar, R. V. (2015) Apoptotic potential of C-phycoerythrin from *Phormidium* sp. A27DM and *Halomicronema* sp. A32DM on human lung carcinoma cells, *EXCLI J.*, **14**: 527–539
- Manirafasha, E., Ndikubwimana T., Zeng , X., Lu, Y., Jing, K. (2016) Phycobiliprotein: Potential microalgae derived pharmaceutical and biological reagent, *Biochemical Engineering Journal*, **109**: 282–296
- Pandey, H. G., Bano, F., Fatma, T. (2011) Studies on *Anabaena* sp. NCCU-9 with special reference to phycocyanin, *J. Algal Biomass Utilization*, **2**(1): 30–51
- Pyo, D., Moon, D. (2005) Adsorption of microcystin LR by activated carbon fibers, *Bulletin of the Korean Chemical Society*, **26** (12):2089-2092

- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C. (1999) Antioxidant activity applying an improved *ABTS* radical cation decolorization assay, *Free Radic Biol Med.*, **26 (9-10)**:1231-7
- Sonani, R. R., Patel, S., Bhastana, B., Jakharia, K., Chaubey, M. G., Singh, N. K., Madamwar, D. (2017) Purification and antioxidant activity of phycocyanin from *Synechococcus* sp. R42DM isolated from industrially polluted site, *Bioresource Technology*, **245** (Pt A):325-331
- Viskari, P. J., Colyer C. L. (2003) Rapid extraction of phycobiliproteins from cultured cyanobacteria samples, *Anal Biochem.* **319(2)**:263-71
- Yan, H., Gong, A., He, H., Zhou, J., Wei, Y., Lv, L. (2006) Adsorption of microcystins by carbon nanotubes, *Chemosphere*, **62 (1)**:142-148

Terrestrial isopods (Isopoda, Crustacea) from the “Danube Delta” Biosphere Reserve

Nicolae Tomescu^{1,✉} and Lucian Alexandru Teodor¹

SUMMARY. The authors describe the fauna of terrestrial isopods and the habitats analysed in 13 sites located in the Biosphere Reserve “Danube Delta”: Letea Forest, Periprava Village, the Levees Maliuc, Caraorman, Dunavăț, Sfântu Gheorghe, Sacalin Island, Popina Island, Enisala Fortress, Sălcișoara (6 Martie) Forest, Doloșman Hill, Gura Portiței and Lupilor Levee. In the investigated habitats 14 species of terrestrial isopods were identified: *Hyloniscus riparius*, *Haplophthalmus orientalis*, *Cylisticus convexus*, *Porcellionides* (= *Metoponorthus*) *pruinus*, *Orthometopon romanicus* n. sp., *Protracheoniscus politus*, *Porcellium collicola*, *Trachelipus arcuatus*, *Trachelipus nodulosus*, *Trachelipus rathkii*, *Trachelipus ratzeburgi*, *Trachelipus squamuliger*, *Armadillidium vulgare*, *Armadillidium jaqueti*.

Keywords: Danube Delta, terrestrial isopods.

Introduction

Research to investigate the flora and fauna of the “Danube Delta” Biosphere Reserve was conducted in the period 1991-1994 under the supervision of the biologist Dr. Vasile Oțel from the Natural Sciences Museum “Danube Delta”, in Tulcea, Tulcea County. Researchers from Bucharest, Cluj, Iași and Constanța participated in this study. Tomescu N. collected terrestrial isopod samples from 13 sites located in the “Danube Delta” Biosphere Reserve: Letea Forest, Periprava Village, the levees Maliuc, Caraorman, Dunavăț, Sfântu Gheorghe, Sacalin Island, Popina Island, Enisala Fortress, Sălcișoara (6 Martie) Forest, Doloșman Hill, Gura Portiței and Lupilor Levee (Fig. 1).

¹ Babeș-Bolyai University, Faculty of Biology and Geology, Department of Taxonomy and Ecology, Cluj-Napoca, Romania.

✉ **Corresponding author:** Nicolae Tomescu, Babeș-Bolyai University of Cluj-Napoca, Department of Taxonomy and Ecology, 5-7 Clinicilor Str., 400006, Cluj-Napoca, Romania.
E-mail: nicolaetomescu36@gmail.com




Figure 1. Map of the study sites investigated in the years 1991-1994, located in the Biosphere Reserve “Danube Delta”: 1 – Letea Forest, 2 – Periprava Village, 3 –Maliuc Levee, 4 – Caraorman Levee, 5 – Dunavăț Levee, 6 – Sântu Gheorghe Levee, 7 – Sacalin Island, 8 – Popina Island, 9 – Enisala Fortress, 10 – Sălchioara (6 Martie) Forest, 11 – Dolosman Hill, 12 – Gura Portiței, 13 – Lupilor Levee.

Fourteen isopod species were identified in the investigated sites, covering a wide range of ecological conditions (habitat types and microhabitats). The species were identified using specific literature: Radu 1983, 1985, Schmalzfuss 1993, 2003, Schmidt 1997, Tomescu 1992, Tomescu *et al.* 2015, Tomescu and Teodor 2016, Vandel 1962, Verhoeff 1907, Wächtler 1937.

Giurginca and Curčić (2003) and Tăbăcaru and Boghean (1989) published terrestrial isopod species lists from Dobrogea, where Popina Island and Enisala are the only locations from the “Danube Delta” Biosphere Reserve mentioned. Tomescu (1992) published terrestrial isopod species from the Caraorman Levee.

The investigated habitats are particularly rich in microhabitats, offering a large variety of environmental conditions for isopods with different ecological requirements (paludal, forest and grassland species).

Materials and methods

Qualitative samples were collected from the 13 sites mentioned in the introduction. Samples were collected directly with tweezers and by using a leaf litter sieve. Isopods were preserved in 70% alcohol. Identification of species followed morphological identification keys from the specialized literature.

The number of samples collected and number of habitats and microhabitats differ in the 13 investigated sites, so a quantitative analysis is impossible to perform. Estimates of the terrestrial isopod fauna were based only on qualitative information. The most frequently sampled habitats in 1991 and 1992 were those present on the Caraorman and Maliuc levees. The other sites were investigated in only one year and over short periods of time, of 1-2 days, following the program established by the project director, biologist dr. Vasile Oțel, who provided transportation of the researchers to the study sites in the Danube Delta.

Description of study site habitats

Letea Forest is dominated by species of poplars, willows and shrubs, growing mostly on dry soil, which is locally interrupted by patches of humid soil on uneven terrain. Thirteen samples from under litter and tree bark from fallen trees were collected in 1993.

Periprava Village is located in the vicinity of Letea forest and has alternating patches of humid soil with rush and dry soils with rare willows and grasslands. The soil of the Danube bank is covered in plant detritus and wood deposits by the houses. Four samples were collected from this area in 1993.

Maliuc Levee. Seven samples were collected from the banks of the Sulina channel, in 1991 and 1992, from habitats dominated by poplars and willows. Isopods were sieved with a leaf litter sieve and collected under rocks and fallen trees.

Caraorman Levee. Caraorman village is surrounded by forests dominated by poplar and willow, oak and ash, or poplar and ash, with forest glades, as well as by grasslands and marshes and rush- covered channel banks. This study site is characterised by a high diversity of habitats and microhabitats. Twenty-five samples were collected from this site in the years 1991-1992.

Dunavăț Levee. Eight litter samples were collected in 1994 from the following habitats: a willow forest with humid soil, under fallen tree trunks, on grasslands, under rocks and around an abandoned building.

Sfântu Gheorghe Levee. Nine samples were collected in 1994 from the following habitats: a poplar forest, an alder forest at the outskirts of Sfântu Gheorghe locality, humid soil flood plains with sedges and rush and channel banks with grassy vegetation.

Sacalin Island. Three samples were collected in 1994 from very humid sandy soil, dominated by rush and sedges and from the edges of rush patches, with less humid soil and with plant detritus.

Popina Island. Six samples were collected in 1992 from grasslands with moderate soil humidity and from the lake shore.

Enisala Fortress. One sample was collected in 1992, from a dry, rocky-soil grassland located at the base of the fortress.

Sălcioara (6 Martie) Forest is mostly composed of oak trees and glades and a pine plantation with humid soil. Five samples were collected here in 1992.

Doloșman Hill. Six samples were collected in 1992 from areas with grassy vegetation, dry soil and lake shore with very humid soil.

Gura Portiței. Two soil samples were collected in 1994 from areas with sandy, highly humid soil, covered with rush and a thick plant detritus layer, and from areas without rush and with moderate soil humidity.

Lupilor Levee. Three samples were collected in 1994 from areas with sandy, highly humid soil, covered with rush and Sea Buckthorn, and from areas without rush, with dry soil and plant detritus.

Results and discussion

Terrestrial isopod species from Letea Forest

Five terrestrial isopod species were identified in the samples collected from Letea Forest: *Hyloniscus riparius*, *Trachelipus arcuatus*, *Trachelipus rathkii*, *Trachelipus ratzeburgi* and *Armadillidium vulgare* (Table 1). The diversity of the investigated microhabitats is reflected by the diversity of the ecological preferences of the species: hygrophilous species (*H. riparius*), forest species (species from the genus *Trachelipus*) and grassland species (*A. vulgare*). *H. riparius*, *T. rathkii* and *A. vulgare* were present in large populations.

Terrestrial isopod species from Periprava Village

Five terrestrial isopod species were found in Periprava Village too: *Hyloniscus riparius*, *Porcellionides* (= *Metoponorthus*) *pruinus*, *Trachelipus rathkii*, *Armadillidium vulgare*, *Armadillidium jaqueti* (Table 1).

Table 1.

Terrestrial isopod species that were identified in the habitats of the “Danube Delta” Biosphere Reserve

Species/ Sites	1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Hyloniscus riparius</i> C. L. Koch, 1838	+	+	-	+	+	+	-	-	-	-	+	-	-
<i>Haplophthalmus orientalis</i> Radu Gh. V., Radu V. V., Cădariu M., 1955	-	-	-	+	-	-	-	-	-	-	+	-	-
<i>Cylisticus convexus</i> De Geer, 1778	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Porcellionides</i> (= <i>Metoponorthus</i>) <i>pruinus</i> Brandt, 1833	-	+	+	+	-	-	-	+	+	-	+	-	-
<i>Orthometopon romanicus</i> Tomescu, Teodor, 2016	-	-	-	-	-	-	+	-	-	-	-	+	+
<i>Protracheoniscus politus</i> C. L. Koch, 1841	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Porcellium collicola</i> Verhoeff, 1907	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Trachelipus arcuatus</i> Budde- Lund, 1885	+	-	+	+	-	-	-	-	-	-	-	-	-
<i>Trachelipus nodulosus</i> C. L. Koch, 1838	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Trachelipus rathkii</i> Brandt, 1833	+	+	+	+	+	+	+	-	-	-	-	+	+
<i>Trachelipus ratzeburg</i> Brandt, 1833	+	-	-	-	-	-	-	-	-	-	-	-	-

Species/ Sites	1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Trachelipus squamuliger</i> Verhoeff, 1907	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Armadillidium vulgare</i> Latreille, 1804	+	+	-	+	+	+	-	-	+	+	+	-	+
<i>Armadillidium jaqueti</i> Dollfus, 1897	-	+	+	-	+	-	-	+	-	-	-	-	-
Total species	5	5	4	7	4	3	2	3	3	4	4	2	3

Stuied sites: **1** – Letea Forest, **2** – Periprava Village, **3** – Maliuc Levee, **4** – Caraorman Levee, **5** – Dunavăț, **6** – Sântu Gheorghe Levee, **7** – Sacalin Island, **8** – Popina Island, **9** – Enisala Fortress, **10** – Sălcioara (6 Martie) Forest, **11** – Doloșman Hill, **12** – Gura Portiței, **13** – Lupilor Levee.

Terrestrial isopod species from the Maliuc Levee

Four species of terrestrial isopods were found in the samples from Maliuc Levee: *Porcellionides pruinosus*, *Trachelipus arcuatus*, *T. rathkii* and *Armadillidium jaqueti*. *T. arcuatus* and *A. jaqueti* (Table 1) were found in relatively large populations.

Terrestrial isopod species from the Caraorman Levee

Seven terrestrial isopod species were identified in the habitats and microhabitats of the Caraorman Levee, all with different ecological requirements: *Hyloniscus riparius*, *Haphlophthalmus orientalis*, *Clysticus convexus*, *Porcellionides pruinosus*, *Trachelipus arcuatus*, *T. rathkii* and *Armadillidium vulgare* (Table 1). In the dry soil of the Caraorman forests we found numerous individuals of *Armadillidium vulgare*, usually a grassland species. Probably, the higher temperatures in the Delta forests favour the dispersal of *A. vulgare* in this habitat type. We found here large populations of the following species: *Trachelipus arcuatus*, *T. rathkii* and *A. vulgare*.

Terrestrial isopod species from the Dunavăț Levee

Four terrestrial isopod species were found in the samples collected from the Dunavăț Levee: *Hyloniscus riparius*, *Trachelipus rathkii*, *Armadillidium vulgare* and *A. jaqueti* (Table 1). The following species had large populations: *Trachelipus rathkii* and *Armadillidium vulgare*.

Terrestrial isopod species from the Sfântu Gheorghe Levee

Three terrestrial isopod species were identified in the samples collected from the Sfântu Gheorghe Levee: *Hyloniscus riparius*, *Trachelipus rathkii* and *Armadillidium vulgare* (Table 1). We found here large populations of the following species: *Trachelipus rathkii* and *A. vulgare*.

Terrestrial isopod species from the Sacalin Island

Two terrestrial isopod species were identified from the samples collected on Sacalin Island (Table 1): *Orthometopon romanicus* n. sp., collected only from very humid soil patches covered with rush and sedges, *Trachelipus rathkii*, collected from areas covered with plant detritus. *O. romanicus* had large populations, with hundreds of individuals per square meter.

Terrestrial isopod species from the Popina Island

Three terrestrial isopod species were identified from the samples collected on Popina Island: *Porcellionides pruinosus*, *Trachelipus nodulosus* and *Armadillidium jaqueti* (Table 1). *P. pruinosus* and *T. nodulosus* had large populations.

Terrestrial isopod species from the Enisala Fortress

Three terrestrial, grassland isopod species were found in the samples collected at Enisala Fortress: *Porcellionides pruinosus*, *Trachelipus nodulosus* and *Armadillidium vulgare* (Table 1). All three species had small populations.

Terrestrial isopod species from the Sălcioara (6 Martie) Forest

Four terrestrial isopod species were identified in the samples collected in the Sălcioara (6 Martie) Forest: *Protracheoniscus politus*, *Porcellium collicola*, *Trachelipus squamuliger* (newly mentioned species for Romania's fauna by Tomescu et al. 2015) and *Armadillidium vulgare* (Table 1). *P. collicola* and *T. squamuliger* had large populations.

Terrestrial isopod species from the Doloşman Hill

Four terrestrial isopod species were identified in the samples collected from the Doloşman Hill: *Hyloniscus riparius*, *Haplophtalmus orientalis*, *Porcellionides pruinosus* and *Armadillidium vulgare* (Table 1).

Terrestrial isopod species from Gura Portiţei

Two terrestrial isopod species were identified in the samples collected from Gura Portiţei: *Orthometopon romanicus* and *Trachelipus rathkii* (Table 1). *O. romanicus* lives in microhabitats that are similar to those from Sacalin Island. Here too, its population counts hundreds of individuals per square meter.

Terrestrial isopod species from Lupilor Levee

Three terrestrial isopod species were identified in the samples collected from Lupilor Levee: *Orthometopon romanicus*, in habitats similar to those on Sacalin Island and Gura Portiței, *Trachelipus rathkii* and *Armadillidium vulgare* (Table 1).

The researches in continental Dobruja mentioned more terrestrial isopod species. Tăbăcaru and Boghean (1989) mentioned 30 species, Giurginca and Ćurčić (2003) mentioned 41 species. Thus, terrestrial isopod communities from continental Dobruja are more ecological diverse than the Danube Delta ones (Table 1).

Conclusions

Fourteen isopod species were identified in the 13 investigated study sites, located in the Biosphere Reserve “Danube Delta”, of which one, *Orthometopon romanicus*, has been described as new for science by Tomescu and Teodor in 2016.

Species widely spread in the investigated sites were: *Hyloniscus riparius*, *Porcellionides* (= *Metoponorthus*) *pruinus*, *Trachelipus rathkii* and *Armadillidium vulgare*.

Species with a limited spread in the investigated sites were: *Haplophthalmus orientalis*, *Cylisticus convexus*, *Porcellium collicola*, *Trachelipus nodulosus*, *T. ratzeburgi*, *T. squamuliger* and *Orthometopon romanicus*.

The number of terrestrial isopod species varies in the different investigated sites according to the number of habitat types and microhabitats, and with the ecological requirements of the species. A relatively large number of species were identified on the Caraorman Levee – seven species – Letea Forest and Periprava Village - each with five species – Maliuc and Dunavăț levees, Sălcioara Forest and Doloșman Hill– each with four species.

The species: *Hyloniscus riparius*, *Orthometopon romanicus*, *Porcellionides pruinus*, *Trachelipus rathkii* and *Armadillidium vulgare* had large populations in the habitats they inhabit.

REFERENCES

- Giurginca, A., Ćurčić, B. S. (2003) A check-list of Oniscidea (Isopoda, Crustacea) from Dobruja (Romania), *Arch. Biol. Sci., Belgrade*, **55**(1-2), 39-44
- Radu, G. V. (1983) *Crustacea, Isopoda, Oniscoidea, Oniscidae inferioare* [in Romanian], Fauna R.S.R., **IV**(13), pp. 168

- Radu, G. V. (1985) *Crustacea, Isopoda, Oniscoidea, Crinocheta* [in Romanian], Fauna R.S.R., **IV**(14), pp. 155
- Schmalfuss, H. (1993) Die Land-Isopoden (Oniscoidea) Griechenlands. 13. Beitrag: Gattung *Orthometopon* („Trachelipidae“) [in German], *Stuttgarter Beitr. Naturk. Ser. A*, **498**, pp. 44
- Schmalfuss, H. (2003) World catalog of terrestrial isopods (Isopoda: Oniscoidea) [in German], *Stuttgarter Beiträge zur Naturkunde*, Serie A. 654, pp. 341
- Schmidt, C. (1997) Revision of the European species of the genus *Trachelipus* Budde-Lund, 1908 (Crustacea: Isopoda: Oniscoidea), *Zoological Journal of Linnean Society*, **121**, 129-244
- Tăbăcaru, I., Boghean, V. (1989) Découverte, en Dobrogea (Roumanie), d'une espèce troglobie du genre *Trachelipus* (Isopoda, Oniscoidea, Trachelipidae) [in French], *Misc. Speol. Rom.*, **1**, 53-75
- Tomescu, N. (1992) Izopode terestre (Crustacea, Izopoda) din Delta Dunării [in Romanian], *An. Ști. Inst. Delta Dunării*, Tulcea, 89-90
- Tomescu N., Teodor L. A., Ferenți S., Covaciu-Marcov S. D. (2015) *Trachelipus* species (Crustacea, Isopoda, Oniscoidea) in Romanian fauna: morphology, ecology, and geographic distribution, *North-Western Journal of Zoology*, **11** (Supplement 1): 1-106
- Tomescu N., Teodor L. A. (2016) *Orthometopon romanicus* n. sp. (Crustacea, Isopoda, Crinocheta: Agnaridae) in Romanian fauna, *Studia UBB Biologia*, **61**(2): 63-68
- Vandel, A. (1962) Isopodes terrestres (Deuxième Partie) [in French], *Faune de France*, **66**, Ed. Paul Lechevalier, Paris, 582-592
- Verhoeff, K. (1907) Über Isopoden. 10. Aufsatz: Zur Kenntnis der Porcellioniden (Körnerasseln) [in German], *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin*, **8**, 229-281
- Wächtler, W. (1937) Ordnung: Isopoda, Asseln [in German], *Die Tierwelt Mitteleuropas II*, Leipzig, **2**, 288-293

Ornithological observations on the Golești Basin (Argeș River, Romania) between February 2013 and January 2014

Adrian Mestecăneanu^{1,✉} and Radu Gava²

SUMMARY. The results of the researches performed between February 2013 and January 2014 on the birds of the Golești reservoir from ROSPA0062 Lacurile de acumulare de pe Argeș are shown in this paper. The 91 observed species belong to 13 orders, Passeriformes being the richest (with 31 species). 53 species are totally or partially dependent on wetlands. The monthly variation of the number of species and individuals reflects both natural and artificial processes. A few species (*Anas platyrhynchos*, *Aythya fuligula*, *Aythya ferina*, *Larus ridibundus*) were noticeable because of their frequency and abundance and this is why they dictated the general dynamics of the local avifauna. As a result, the Anseriformes and Charadriiformes are the overdominant orders at general level. 21 species are included in the Annex I of the Birds Directive. The qualitative and quantitative alterations observed over time do not show necessarily the climatic changes, because the conditions of the basin did not remain constant.

Keywords: anthropogenic pressure, avifauna, basin, Special Protected Area.

Introduction

The avifauna of the reservoirs from Romania has been the subject of many works since the construction of the artificial lakes. In these papers the influence of the latter on the birds was constantly highlighted (Munteanu, 1978, Munteanu, 2000, Gache, 2002, Mitruțy, 2002, Rang, 2002 etc.).

The first thematic study about the middle and upper hydrographic basin of the Argeș River was performed before the construction of the reservoirs (Mătieș, 1969). Their own avifauna was subsequently studied, when their major importance as places of stopover or wintering was revealed (Munteanu and Mătieș, 1983). Other data about them was published later, in a synthesis work on the aquatic birds from the winter quarters from Romania (Munteanu *et al.*, 1989) and in 1997 it was suggested

¹ Argeș County Museum, Armand Călinescu Str., No. 44, 110047, Pitești, Argeș, Romania.

² University of Pitești, Târgu din Vale Str., No. 1, 110040, Pitești, Argeș, Romania.

✉ **Corresponding author: Adrian Mestecăneanu**, Argeș County Museum, Armand Călinescu Str., No. 44, 110047, Pitești, Argeș, Romania,
E-mail: mestecaneanua@yahoo.com

the possibility for some of these locations to become Important Bird Area (Gava, 1997). The research about the reservoirs' ornithofauna between Vâlcele and Golești has been intensified after 2004 (Gava *et al.*, 2004a,b, 2007, 2008, 2011, 2012, Mestecăneanu *et al.*, 2004, 2006b, 2008, 2010, 2013, Conete *et al.*, 2006, 2008, 2010, 2011, Mestecăneanu and Gava, 2013, 2015a,b, 2016a,b,c, 2017 etc.) and, in this context, some papers were dedicated to the birds from the Golești Basin (Mestecăneanu *et al.*, 2005, 2006a, Conete *et al.*, 2009, 2012). Also, a PhD Thesis referred to the birds from the area (Conete, 2011).

Materials and methods

The Golești Basin is a component of the protected site ROSPA0062 Lacurile de acumulare de pe Argeș ("The Basins from the Argeș River"), included in Natura 2000 Network. It is the southern element of this series of basins that begins at Zigoneni, i.e. upstream, and continues downstream with the reservoirs Vâlcele, Budeasa, Bascov and Pitești (Fig. 1). Its features are: type – gravity dam/earth; kind of sealing – uphill embankment, concrete; height – 32 m; length – 7,866 m; volume – $78.5 \times 10^6 \text{ m}^3$; area – 680 ha; length of the lake – 7 km; object – water supply, electricity, flood prevention, irrigation; area of catchment – 3100 km²; discharge flow – 3,760 m³/s; spill type – overflowing with the edges; gift in use - 1983 (cf. <http://www.baraje.ro>).




Figure 1. The map of the area with the place of the Golești Basin, marked with blue arrow.

The climate of the area is temperate-continental, with traits of plain. The average annual temperature of the air is 10°C; in January, it is 2.7 °C and in July it is nearly 21 °C (Măciu *et al.*, 1982). The average annual temperature of the water is bigger with 1-3 °C than the one of the air and it is ca. 9 °C at Pitești, 8 km far away from Golești. In the harsh winters, the ice bridge is formed in the first half of January and disappears in the last part of February (Barco and Nedelcu, 1974).

The vegetation is composed by species of the genera *Carex*, *Juncus*, *Phragmites* that, together with a few mature alders, grow toward the end of the lake where the right bank is covered with grass and other herbaceous species: *Agrimonia eupatoria* L., *Carex sylvatica* Huds., *Urtica dioica* L., *Lysimachia nummularia* L. and woody vegetation: *Alnus glutinosa* (L.), *Prunus spinosa* L., *Crataegus monogyna* Jacq., *Rubus caesius* L. etc. The floating vegetation emerges mainly in the upstream extremity: *Myriophyllum verticillatum* L., *Sparganium erectum* L., *Mentha aquatica* L., *Polygonum mite* L., *Lemna minor* L. etc. Toward the banks, *Typha* sp. can be found (Stancu, 2014).

Between the water surface and the forest of *Quercus* from the right bank, there is a bare band of gravel. Except the treed upstream end, the left bank and the dam have no vegetation because of the concrete bevel.

The area from vicinity is cultivated mainly with cereal crops, cabbage and grass.

The fish fauna is diverse: *Alburnus alburnus* Linnaeus, 1758, *Barbus barbus* (Linnaeus, 1758), *Barbus meridionalis petenyi* (Heckel, 1847), *Carassius auratus gibelio* (Bloch, 1782), *Gobio gobio* (Linnaeus, 1758), *Squalius cephalus* (Linnaeus, 1758), *Rutilus rutilus* (Linnaeus, 1758), *Rhodeus amarus* (Bloch, 1782), *Pseudorasbora parva* (Temminck & Schlegel, 1846), *Cobitis taenia* Linnaeus, 1758, *Esox lucius* Linnaeus, 1758, *Perca fluviatilis* Linnaeus, 1758, *Lepomis gibbosus* (Linnaeus, 1758), *Sander lucioperca* (Linnaeus, 1758) (Truță and Dumitru, 2015). Many of the species represent the food source for the ichthyophagous birds.

There is a restricted road to the public circulation on the bevel of the dam. A motorway goes alongside the right end of the basin. Toward South (Recea and Cătanele), East (Udeni-Zăvoi) and West (Pitești and Bradu) there are human settlements, situated under 500 m distance (Fig. 2).

The itinerary method combined to the one of fixed points of observations was used. Monthly, one field trip was performed, generally between the 10-th and 20-th day. Our main goal was to count the birds from the water surface and, in order to do that, we walked on the bevel – a place with good visibility. The amount of the birds found in big number was two times estimated, resulting mean values, and the strengths of the weakly represented species were precisely numbered. A particular care was to avoid as possible the double counting. The birds were visually and auditory identified. Two binoculars (10x50), a spotting scope (14-45x50) and a camera (42x optical zoom) were used.

The scientific norm and classification of the birds are compatible with the Hamlin Guide (Bruun *et al.*, 1999).



Figure 2. Aerial view of the Golești Basin, by Google Earth.

Results and discussion

During the above mentioned interval of time, 91 species of birds were identified. They represent less than a half of the 199 species recorded here during the preceding study (Conete *et al.*, 2012) and this is mainly explained by the fact the period of monitoring was much shorter (a year versus eight years). So, if we eliminate the 86 subprecedent species from the Dzuba index of ecological significance point of view (that means rare and very rare species, represented by few individuals) previously recorded, we obtain a figure which is closer to the one registered now.

The species currently observed belong to 13 orders (Gaviiformes – 1 species, Podicipediformes – 4 species, Pelecaniformes – 3 species, Ciconiiformes – 7 species, Anseriformes – 13 species, Falconiformes – 6 species, Galliformes – 2 species, Gruiformes – 2 species, Charadriiformes – 20 species, Cuculiformes – 1 species, Apodiformes – 1 species, Coraciiformes – 1 species, Piciformes – 1 species and Passeriformes, the most numerous – 31 species), 4 (Columbiformes, Strigiformes, Caprimulgiformes and Piciformes) less than during 2003-2010.

53 species are totally or partially dependent on wetlands (Table 1). They belong to 9 orders: Gaviiformes – 1 species, Podicipediformes – 4 species, Pelecaniformes – 3 species, Ciconiiformes – 7 species, Anseriformes – 13 species, Falconiformes – 1 species, Gruiformes – 2 species, Charadriiformes, the best represented – 20 species, and Passeriformes – 2 species.

The number of species was the biggest in April (35) and August (40), corresponding to the migration periods (Table 2). A relatively big number was recorded in May – July (between 26 and 30 species), in the breeding period for the most of the species, while in the hiemal season, their number was, generally the

lowest (between 22 and 28). For the species dependent on wetlands, the principal maximum was in February (18 species) and the secondary one, in August (15 species); in April was registered the third peak (10 species). In the breeding period, the number of species was generally low (minimum 4 species, in May), while in the hiemal season the number of species increased from November (5 species) to February (18 species, as previously shown).

The 45,961 individuals counted along the year had a different dynamics (Table 2). The biggest strength was noticed in February (8,372 individuals) while lower peaks were recorded in August (7,505 individuals) and November (7,209 individuals). The maximum is over 45% of the one registered on all the basins from the Argeș River, upstream Pitești, before 1980 (Munteanu and Mătieș, 1983), a number which is very close to that recorded currently only on five basins from the segment of the Argeș River between Vâlcele and Golești (Mestecăneanu and Gava, 2016a). The lowest values were between April and June (the minimum, in May – 184 individuals), but a small number was also registered in December (1,254 individuals). As regard the strength of the species dependent on wetlands, which totalised 44,078 individuals, the dynamics varied identically, but the values were lower than the firsts with 69 (in December) to 382 individuals (in the passage from September), that show that the input of the non-wetland species was small (it must be more significant, our attention being focused on the other group of species). The scarcity of individuals from April to June was also noticed on all the basins from the upper and middle course of the Argeș River until 1980, when 84 aquatic species were counted (Munteanu and Mătieș, 1983).

These reflect the migration periods, mainly for the birds dependent on wetlands, and, also, show that the Golești Basin is a very attractive place of wintering, when a moderate number of species provides a relative big number of individuals. As well as over 30 years ago (Munteanu and Mătieș, 1983), even now the basin is not good for breeding, because of the limited perimeter of aquatic and amphibious vegetation, but, in the future the situation is expected to become more favourable as a consequence of the developing of the natural silting and afforesting processes. Beside the intrinsic factors (the fluctuation of the water level, the variation of the food supply and shelters, in the cold time in relation with the gradual freezing or thawing of the water surface, that determine the migration etc.), the local dynamics of the avifauna can be negatively influenced by the anthropogenic elements (fishing, hunting, pasturage and other intrusions), as we discussed before (Mestecăneanu and Gava, 2015b, 2016a, c) but also positively, because large interventions on the surrounding dam lakes (the complete desiccations or the presence of many boats on the water, for instance) determine many birds to move, and thus to increase temporarily the number of individuals and species from here.

A correlation with the surface covered with ice cannot be made, because this was reduced (ca. 5%) and it was formed only in January. Instead, a simple correlation between the number of species dependent on wetland and the number of

fishermen can be obtained: it is -0.14 (negative and weak correlation) while the one between the number of individuals for the same species and the number of fishermen is -0.35 (negative and acceptable correlation, by Colton, 1974). That means that the fishermen affect to a certain extent the presence of the birds in the area, even if they were observed both on banks and on boats only in 75% of samples, with a maximum density (in April and October) of 0.04 persons/ha.

Table 1.

The occurrence along the year and some ecological indexes.

No.	Species	January	February	March	April	May	June	July	August	September	October	November	December	Absolute abundance	Class of constancy	Class of dominance	Class of Dzuba index
I. Gaviiformes																	
1	<i>Gavia arctica</i> (Linnaeus, 1758)*												+	1	C1	D1	W1
II. Podicipediformes																	
2	<i>Podiceps cristatus</i> (Linnaeus, 1758)*	+	+	+	+	+	+	+	+	+	+	+	+	760	C4	D2	W3
3	<i>Podiceps grisegena</i> Boddaert, 1783*								+					2	C1	D1	W1
4	<i>Podiceps nigricollis</i> Brehm, 1831*				+									2	C1	D1	W1
5	<i>Tachybaptus ruficollis</i> (Pallas, 1764)*		+								+			13	C1	D1	W1
III. Pelecaniformes																	
6	<i>Phalacrocorax carbo</i> (Linnaeus, 1758)*	+		+		+	+	+	+	+	+	+	+	301	C4	D1	W2
7	<i>Phalacrocorax pygmeus</i> (Pallas, 1773)*	+		+								+	+	52	C2	D1	W1
8	<i>Pelecanus crispus</i> Bruch, 1832*						+		+					13	C1	D1	W1
IV. Ciconiiformes																	
9	<i>Ixobrychus minutus</i> (Linnaeus, 1766)*								+					1	C1	D1	W1
10	<i>Egretta garzetta</i> (Linnaeus, 1766)*				+	+	+	+	+					90	C2	D1	W1
11	<i>Egretta alba</i> (Linnaeus, 1758)*		+		+					+				4	C1	D1	W1
12	<i>Ardeola ralloides</i> (Scopoli, 1769)*					+								1	C1	D1	W1
13	<i>Ardea cinerea</i> Linnaeus, 1758*	+	+		+			+	+	+	+			79	C3	D1	W2

ORNITHOLOGICAL OBSERVATIONS ON THE GOLEȘTI BASIN

No.	Species	January	February	March	April	May	June	July	August	September	October	November	December	Absolute abundance	Class of constancy	Class of dominance	Class of Dzuba index
14	<i>Nycticorax nycticorax</i> (Linnaeus, 1758)*					+	+	+						13	C1	D1	W1
15	<i>Ciconia ciconia</i> (Linnaeus, 1758)*						+	+						2	C1	D1	W1
V. Anseriformes																	
16	<i>Cygnus olor</i> (Gmelin, 1789)*	+	+	+	+	+	+	+	+	+		+	+	173	C4	D1	W2
17	<i>Cygnus cygnus</i> (Linnaeus, 1758)*		+											9	C1	D1	W1
18	<i>Anser albifrons</i> (Scopoli, 1769)*	+											+	660	C1	D2	W2
19	<i>Anas platyrhynchos</i> Linnaeus, 1758*	+	+	+	+	+	+	+	+	+	+	+	+	13812	C4	D5	W5
20	<i>Anas penelope</i> Linnaeus, 1758*		+	+									+	50	C1	D1	W1
21	<i>Anas querquedula</i> Linnaeus, 1758*				+				+	+				278	C1	D1	W2
22	<i>Anas crecca</i> Linnaeus, 1758*	+	+	+	+				+	+	+	+	+	20023	C3	D3	W3
23	<i>Anas chryseola</i> Linnaeus, 1758*						+		+					19	C1	D1	W1
24	<i>Tadorna tadorna</i> (Linnaeus, 1758)*		+						+				+	49	C1	D1	W1
25	<i>Aythya fuligula</i> (Linnaeus, 1758)*	+	+	+	+			+	+	+	+	+	+	5209	C3	D5	W4
26	<i>Aythya ferina</i> (Linnaeus, 1758)*	+	+	+	+		+	+	+	+	+	+	+	10770	C4	D5	W5
27	<i>Aythya nyroca</i> Gldenstdt, 1770*							+						2	C1	D1	W1
28	<i>Bucephala clangula</i> (Linnaeus, 1758)*	+	+	+									+	252	C2	D1	W2
VI. Falconiformes																	
29	<i>Buteo buteo</i> (Linnaeus, 1758)	+	+		+							+	+	19	C2	D1	W1
30	<i>Circus aeruginosus</i> (Linnaeus, 1758)*				+									3	C1	D1	W1
31	<i>Falco subbuteo</i> Linnaeus, 1758						+							1	C1	D1	W1
32	<i>Falco vespertinus</i> Linnaeus, 1766									+				1	C1	D1	W1
33	<i>Falco tinnunculus</i> Linnaeus, 1758			+	+	+		+	+			+		11	C2	D1	W1
VII. Galliformes																	
34	<i>Phasianus colchicus</i> Linnaeus, 1758	+	+									+		14	C1	D1	W1
35	<i>Coturnix coturnix</i> (Linnaeus, 1758)					+								1	C1	D1	W1

No.	Species													Absolute abundance	Class of constancy	Class of dominance	Class of Dzuba index		
		January	February	March	April	May	June	July	August	September	October	November	December						
VIII. Gruiformes																			
36	<i>Gallinula chloropus</i> (Linnaeus, 1758)*					+										1	C1	D1	W1
37	<i>Fulica atra</i> Linnaeus, 1758*	+	+	+	+	+	+	+		+	+	+	+			9831	C4	D3	W3
IX. Charadriiformes																			
38	<i>Vanellus vanellus</i> (Linnaeus, 1758)*			+			+		+	+						39	C2	D1	W1
39	<i>Charadrius dubius</i> Scopoli, 1786*				+				+							8	C1	D1	W1
40	<i>Gallinago gallinago</i> (Linnaeus, 1758)*								+							1	C1	D1	W1
41	<i>Limosa limosa</i> (Linnaeus, 1758)*								+							1	C1	D1	W1
42	<i>Calidris alpina</i> (Linnaeus, 1758)*								+							1	C1	D1	W1
43	<i>Calidris minuta</i> (Leisler, 1812)*								+							2	C1	D1	W1
44	<i>Actitis hypoleucos</i> (Linnaeus, 1758)*							+	+							25	C1	D1	W1
45	<i>Tringa ochropus</i> Linnaeus, 1758*	+	+		+				+							10	C2	D1	W1
46	<i>Tringa glareola</i> Linnaeus, 1758*								+							22	C1	D1	W1
47	<i>Tringa nebularia</i> (Gunnerus, 1767) *								+							4	C1	D1	W1
48	<i>Philomachus pugnax</i> (Linnaeus, 1758)*								+							4	C1	D1	W1
49	<i>Recurvirostra avosetta</i> Linnaeus, 1758*				+											7	C1	D1	W1
50	<i>Himantopus himantopus</i> (Linnaeus, 1758)*								+							30	C1	D1	W1
51	<i>Larus argentatus</i> Pontoppidan, 1763*	+	+	+	+	+	+	+	+	+	+	+	+	+	+	2236	C4	D2	W3
52	<i>Larus canus</i> Linnaeus, 1758*	+								+		+	+			2025	C2	D2	W2
53	<i>Larus ridibundus</i> Linnaeus, 1766*	+	+	+	+	+	+	+	+	+	+	+	+	+	+	13163	C4	D5	W5
54	<i>Larus minutus</i> Pallas, 1776*								+							4	C1	D1	W1
55	<i>Chlidonias niger</i> (Linnaeus, 1758)*								+							9	C1	D1	W1

ORNITHOLOGICAL OBSERVATIONS ON THE GOLEȘTI BASIN

No.	Species	January	February	March	April	May	June	July	August	September	October	November	December	Absolute abundance	Class of constancy	Class of dominance	Class of Dzuba index
56	<i>Chlidonias hybridus</i> (Pallas, 1811)*						+	+						24	C1	D1	W1
57	<i>Sterna hirundo</i> Linnaeus, 1758*				+	+	+	+						30	C2	D1	W1
X. Cuculiformes																	
58	<i>Cuculus canorus</i> Linnaeus, 1758					+								1	C1	D1	W1
XI. Apodiformes																	
59	<i>Apus apus</i> (Linnaeus, 1758)					+								32	C1	D1	W1
XII. Coraciiformes																	
60	<i>Upupa epops</i> Linnaeus, 1758			+	+									5	C1	D1	W1
XIII. Passeriformes																	
61	<i>Galerida cristata</i> (Linnaeus, 1758)		+							+				6	C1	D1	W1
62	<i>Alauda arvensis</i> Linnaeus, 1758			+	+	+	+							10	C2	D1	W1
63	<i>Riparia riparia</i> (Linnaeus, 1758)					+	+		+					251	C1	D1	W1
64	<i>Hirundo rustica</i> Linnaeus, 1758				+	+	+	+	+	+				204	C2	D1	W2
65	<i>Delichon urbica</i> (Linnaeus, 1758)						+	+						90	C1	D1	W1
66	<i>Anthus trivialis</i> (Linnaeus, 1758)				+									9	C1	D1	W1
67	<i>Anthus spinoletta</i> (Linnaeus, 1758)										+			40	C2	D1	W1
68	<i>Motacilla flava</i> Linnaeus, 1758				+	+	+	+	+					32	C2	D1	W1
69	<i>Motacilla alba</i> Linnaeus, 1758			+	+			+	+	+	+			136	C2	D1	W1
70	<i>Sturnus vulgaris</i> Linnaeus, 1758			+	+	+	+			+	+	+		371	C3	D1	W2
71	<i>Pica pica</i> (Linnaeus, 1758)	+	+	+	+	+	+	+	+	+	+	+	+	391	C4	D1	W2
72	<i>Corvus monedula</i> Linnaeus, 1758	+	+	+		+	+	+	+	+	+	+	+	1260	C4	D1	W2
73	<i>Corvus frugilegus</i> Linnaeus, 1758	+	+	+	+	+	+		+	+	+	+	+	1104	C4	D1	W2

No.	Species	January	February	March	April	May	June	July	August	September	October	November	December	Absolute abundance	Class of constancy	Class of dominance	Class of Dzuba index
74	<i>Corvus corone cornix</i> Linnaeus, 1758	+	+	+		+			+	+		+	+	21	C3	D1	W1
75	<i>Corvus corax</i> Linnaeus, 1758	+	+	+	+							+	+	41	C3	D1	W1
76	<i>Prunella modularis</i> (Linnaeus, 1758)										+			5	C1	D1	W1
77	<i>Acrocephalus palustris</i> Bechstein, 1798*						+							30	C1	D1	W1
78	<i>Sylvia communis</i> Latham, 1787					+								4	C1	D1	W1
79	<i>Phylloscopus collybita</i> Vieillot, 1817								+					17	C1	D1	W1
80	<i>Oenanthe oenanthe</i> (Linnaeus, 1758)				+									6	C1	D1	W1
81	<i>Luscinia luscinia</i> (Linnaeus, 1758)								+					2	C1	D1	W1
82	<i>Remiz pendulinus</i> (Linnaeus, 1758)*							+						8	C1	D1	W1
83	<i>Passer domesticus</i> (Linnaeus, 1758)			+			+			+		+		245	C2	D1	W1
84	<i>Passer montanus</i> (Linnaeus, 1758)						+		+					211	C1	D1	W1
85	<i>Fringilla coelebs</i> Linnaeus, 1758										+			47	C1	D1	W1
86	<i>Carduelis chloris</i> (Linnaeus, 1758)				+						+			35	C1	D1	W1
87	<i>Carduelis spinus</i> (Linnaeus, 1758)											+		7	C1	D1	W1
88	<i>Carduelis carduelis</i> (Linnaeus, 1758)			+		+					+		+	65	C2	D1	W1
89	<i>Carduelis cannabina</i> (Linnaeus, 1758)		+		+							+	+	24	C2	D1	W1
90	<i>Miliaria calandra</i> (Linnaeus, 1758)		+	+	+	+	+				+	+		34	C3	D1	W1
91	<i>Emberiza citrinella</i> Linnaeus, 1758	+									+	+		188	C1	D1	W1

Legend:* - birds that depend on wetlands; + - presence; C1 – accidental species, C2 – accessory species, C3 – constant species, C4 – euconstant species; D1, W1 – subrecedent species, D2, W2 – recedent species, D3, W3 – suddominant species, D4, W4 – dominant species, D5, W5 – eudominant species; AI, AII, AIII – annexes of the Birds Directive, Bern Convention and, respectively, Bonn Convention, A, B – parts of the annexes.

Table 2.

The dynamics of the species and their strengths.

Month	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.
Number of species	24	28	27	35	28	30	26	42	24	26	25	22
Number of individuals	6,163	8,372	2,316	607	184	542	1,582	7505	3,278	6,949	7,209	1,254
Number of species*	8	18	7	10	4	6	6	15	6	6	5	6
Number of individuals*	6,036	8,300	2,189	523	49	354	1,477	7,228	2,896	6,767	7,074	1,185

Legend: * - birds that depend on wetlands.

Referring to the ecological indexes, regarding the constancy, 57 species (62.64%) were occasional (C1), 16 species (17.58%) were accessory (C2), 7 species (7.69%) were constant (C3) and 11 species (12.09%) were euconstant (C4), (Table 1, Fig. 3). The euconstant species were: *Podiceps cristatus*, *Phalacrocorax carbo*, *Cygnus olor*, *Anas platyrhynchos*, *Aythya ferina*, *Fulica atra*, *Larus argentatus*, *Larus ridibundus*, *Pica pica*, *Corvus monedula*, and *Corvus frugilegus*. *Podiceps cristatus*, *Anas platyrhynchos*, *Larus argentatus*, *Larus ridibundus* and *Pica pica* were observed every month, *Cygnus olor*, *Fulica atra*, *Corvus monedula*, and *Corvus frugilegus* were observed 11 times and *Phalacrocorax carbo* and *Aythya ferina* were observed 10 times (Table 1). *Pica pica*, *Corvus monedula*, and *Corvus frugilegus* (all residents, omnivorous, from Corvidae family) are the only non-wetland birds, while the others are birds adapted to the wetlands. *Larus argentatus* was represented by two subspecies: *L. cachinnans* and *L. michahellis* (different species, by other classifications), where the last, which breeds in Pitești, was encountered all year round.




Figure 3. The species distribution according to the index of constancy (C1 – accidental species, C2 – accessory species, C3 – constant species, C4 – euconstant species).

By the dominance (Table 1, Fig. 4), the majority of the species (81) were subrecedent (89.09%). Only 4 (4.40%) were recedent, 2 (2.20%) were subdominant, and 4 (4.40%) were eudominant. It is surprising the absence of the dominant species. The distribution of the assembly suggests a large discrepancy between a big number of species represented by few individuals and a small number of species represented by many individuals, as we saw in other occasion, too (Mestecăneanu and Gava, 2016a). The group of the eudominant species is constituted by three species of Anseriformes (*Anas platyrhynchos*, *Aythya fuligula*, and *Aythya ferina*) and one species of Charadriiformes (*Larus ridibundus*). They are species with a vast range of food items (omnivorous, the first three, and zoofagous-polyfagous, the last one), and except *Aythya fuligula*, mainly winter visitor, the others are preponderantly partial migratory in our country (Bruun *et al.*, 1999). Before 1980, *Anas platyrhynchos* was the most abundant species on the Argeş basins, too; it was followed by *Anas crecca*, *Anas querquedula*, *Vanellus vanellus* and then by *Larus ridibundus* (Munteanu and Mătieş, 1983).




Figure 4. The species distribution according to the index of dominance (D1 – subrecedent species, D2 – recedent species, D3 – subdominant species, D4 – dominant species, D5 – eudominant species).

By the Dzuba index of ecological significance, that takes into consideration both constancy and dominance, the most of the species (72) were subrecedent (89.09%); 11 (12.09%) were recedent, 4 (4.40%) were subdominant, 1 (1.10%, *Aythya fuligula*) was dominant and 3 (3.30%, *Anas platyrhynchos*, *Aythya ferina*, and *Larus ridibundus*) were eudominant (Table 1, Fig. 5).




Figure 5. The species distribution according to the index of Dzuba ecological significance (W1 – subrecedent species, W2 – recedent species, W3 – subdominant species, W4 – dominant species, W5 – eudominant species).

Anas platyrhynchos had the biggest strength in February (4,100 individuals) and November (3,212 individuals), but big numbers were also registered in January (2,100) and October (2,000). The fewest individuals were observed in May (8 individuals) and generally between April and July (below 119). Even if it was a constant presence, the species is only probable breeding in the area. *Aythya ferina* had the biggest strengths in October (2,530 individuals) and February (2,350 individuals). The lowest numbers were registered from March to June, and in September. In May the figure was zero, the species being only possible breeding. Unexpectedly, the strength was null in December, too, when also the general number of all the observed species was low, by comparison to the one recorded in the other months of the hiemal season and on the other basins from the Argeș River at this time, fact that suggests a massive derange just before our visit. *Larus ridibundus* had the maximum strength in August (3,184 individuals) and in January (1,150 individuals). During the rest of the year it was lower, inclusively in December, usually one of the best represented months. Although it was registered in every field trip, it is only a possible breeder in the area (Fig. 6).

Regarding the density of these species (Table 3), we observe that *Anas platyrhynchos* attains 6.06 individuals/ha in February and 4.42 individuals/ha in November, *Aythya ferina* reaches 3.72 individuals/ha in October and 3.46 individuals/ha in February, and *Larus ridibundus* arrives to 4.68 individuals/ha in August and 1.69 individuals/ha, in January. All of them have 9.59 individuals/ha in February and 8.21 individuals/ha in August. In the same time, all the recorded species attain 12.31 individuals/ha in February and 11.04 individuals/ha in August, while the species dependent on wetlands have 12.21 individuals/ha in February, respectively 10.40 individuals/ha in August. Big values also

characterised October and November. Beside the very obvious migration of *Larus ridibundus* noticed in August, the increasing of the ratio stated now can be related to some species of Anseriformes and Charadriiformes: because they find shelter and food, the first ones gather on the basin for moulting from inadequate waters of the area and the second ones halt here in migration. Over 30 years ago, this density had maximum 15 individuals/ha (Munteanu and Mătieș, 1983), but the considered area is not clearly mentioned.




Figure 6. The monthly variation of the strength for the eudominant species from the Dzuba index point of view.

Table 3.

The monthly and yearly number of individuals/ha for some species.

Period	January	February	March	April	May	June	July	August	September	October	November	December	Total
<i>Anas platyrhynchos</i>	3.09	6.03	0.22	0.07	0.01	0.18	0.08	1.18	1.10	2.94	4.72	0.69	20.31
<i>Aythya ferina</i>	1.54	3.46	0.62	0.06	0.00	0.19	1.62	2.35	0.51	3.72	1.76	0.00	15.84
<i>Larus ridibundus</i>	1.69	0.10	0.11	0.04	0.00	0.01	0.06	4.68	0.62	0.08	1.10	0.02	8.52
Eudominant species	6.32	9.59	0.95	0.17	0.01	0.38	1.75	8.21	2.24	6.74	7.59	0.71	44.67
All species	9.06	12.31	3.41	0.89	0.27	0.80	2.33	11.04	4.82	10.22	10.60	1.84	67.59
Wetland species	8.88	12.21	3.22	0.77	0.07	0.52	2.17	10.63	4.26	9.95	10.40	1.74	64.82

By the index of relation (IR), as a result of the specific dominance, Anseriformes and Charadriiformes were the overdominant orders, the first with 72.02% and the second with 15.77%. There were not dominant orders, because the others are complementary (Fig. 7).




Figure 7. The participation of the orders to the formation of the avicoenose, by the index of relation – IR (SA – the static axis, DA – the dominance axis).

The monthly dynamics (Fig. 8) shows Anseriformes permanently placed in the overdominant zone, except May, when it was complementary. Charadriiformes was overdominant in August, September and January, dominant in March, April, November and December, and complementary in the rest of the year. Together, the other orders were overdominant in May, especially due to the Passeriformes, but also in March, April, June, July and September, on the background of the smaller strengths of Anseriformes and Charadriiformes.




Figure 8. The monthly dynamic of the orders, by the IR (SA – the static axis, DA – the dominance axis).

Within the Anseriformes order (Fig. 9), *Anas platyrhynchos*, *Aythya ferina* and *Aythya fuligula* were the overdominant species, while the others are complementary, and within the Charadriiformes order (Fig. 10), *Larus ridibundus* and *Larus argentatus* were the overdominant species, *Larus canus* was the dominant species and the others were complementary.

The monthly dynamics of the species do not reveal continuous overdominant species (Fig. 11, Fig. 12). However, most of the time, *Anas platyrhynchos* and *Aythya ferina*, respectively *Larus ridibundus* and *Larus argentatus*, were in the overdominance zone. *Aythya fuligula* was overdominant in March, April and October, November, and *Larus canus* only in January. Notable is the group of the other species which was also overdominant in April, May, August, September and December (due to the presence of *Anser albifrons*), among the Anseriformes, and between April and July (due to *Recurvirostra avosetta*, *Sterna hirundo*, *Vanellus vanellus*, or *Chlydonias hybridus*), among the Charadriiformes.




Figure 9. The participation of the species to the Anseriformes coenose, by the IR (SA – the static axis, DA – the dominance axis).




Figure 10. The participation of the species to the formation of Charadriiformes coenose, by the IR (SA – the static axis, DA – the dominance axis).

ORNITHOLOGICAL OBSERVATIONS ON THE GOLEȘTI BASIN




Figure 11. The monthly dynamic of the species inside the Anseriformes order, by the IR (SA – the static axis, DA – the dominance axis).




Figure 12. The monthly dynamic of the species inside the Charadriiformes order, by the IR (SA – the static axis, DA – the dominance axis).

Regarding the protection, 21 species (23.07%: *Gavia arctica*, *Phalacrocorax pygmeus*, *Pelecanus crispus*, *Ixobrychus minutus*, *Egretta garzetta*, *Egretta alba*, *Ardeola ralloides*, *Nycticorax nycticorax*, *Ciconia ciconia*, *Cygnus cygnus*, *Aythya nyroca*, *Circus aeruginosus*, *Calidris alpina*, *Tringa glareola*, *Philomachus pugnax*, *Recurvirostra avosetta*, *Himantopus himantopus*, *Larus minutus*, *Chlidonias niger*, *Chlidonias hybridus*, and *Sterna hirundo*) included in the Annex I by the Birds Directive – the Council Directive 2009/147/EC were observed in the perimeter of the basin. All of them are species dependent on wetlands and represent 39.62%.

Measures for the habitat protection to assure their survival and reproduction in their area of distribution must be applied (<http://ec.europa.eu/environment/nature/legislation/>). Among them, *Phalacrocorax pygmeus*, *Egretta garzetta* and *Sterna hirundo* are accessory. *Phalacrocorax pygmeus* was observed mainly in the hiemal season, and *Egretta garzetta* and *Sterna hirundo* were registered in the breeding period; they are possible, respectively probable breeding species on the basin. Regarding the dominance, all the species from the Annex are subprecedent. The most abundant were *Egretta garzetta* (90 individuals) and *Phalacrocorax pygmeus* (52 individuals).

Conclusions

The avifauna observed between February 2013 and January 2014 on the Golești Basin was diverse, with 91 species that belong to 13 orders. Among them, 53 are totally or partially dependent on wetlands.

The dynamics of the species and their strengths shows maximums that correspond to the migration and wintering times. Because of the unsatisfactory conditions, the breeding period was characterised by few species and individuals. The fluctuations reflect both the natural processes and the human pressure, a negative correlation between the fishing and the birds' number being shown.

It seems to be an increase of the number of the individuals in relation to the strengths recorded in the zone of the basins over 30 years ago, but the fact is unclear because of the different method and area of data collecting. Some acts of eutrophication, the installation of vegetation and the climatic changes can be included here.

Several species (*Anas platyrhynchos*, *Aythya fuligula*, *Aythya ferina*, *Larus ridibundus*) were noticed due to their frequency and abundance. They trigger the dynamics of the local avifauna. As result, Anseriformes and Charadriiformes were the overdominant orders. The overdominant species were *Anas platyrhynchos*, *Aythya ferina* and *Aythya fuligula* inside the Anseriformes order, and *Larus ridibundus* and *Larus argentatus* inside the Charadriiformes order.

21 species are included in the Annex I by the Birds Directive; some of them are frequent species during winter (*Phalacrocorax pygmeus*), respectively summer (*Egretta garzetta* and *Sterna hirundo*). All of these and the big number of species and individuals justify the necessity of considering Golești Basin a protected area and the reason to diminish the negative anthropogenic impact as a priority in the future.

REFERENCES

- Bruun, B., Delin, H., Svensson, L., Singer, A., Zetterström, D., Munteanu, D. (1999) *Păsările din România și Europa. Determinator ilustrat*, Hamlyn Guide, Societatea Ornitologică Română, Octopus Publishing Group Ltd., pp. 320
- Colton, T. (1974) *Statistics in Medicine*, Little Brown and Company, New York, pp. 224
- Conete, D. (2011) *Cercetări ecologice asupra avifaunei unor lacuri de baraj din zona mijlocie a văii Argeșului*, PhD thesis, Institutul de Biologie al Academiei Române București, pp. 370
- Conete, D., Gava, R., Mestecăneanu, A. (2008) Statutul de protecție al păsărilor din zona lacurilor de acumulare de pe râul Argeș, *Scripta Ornithologica Romaniae*, Cluj-Napoca, **3**: 68-75
- Conete, D., Gava, R., Mestecăneanu, A. (2009) Ecological researches on the avifauna of the Golești basin in the hiemal and prevernal aspects (2008 - 2009), *Annals, Food Science and Tehnology*, Universitatea Valahia, Târgoviște, **10**, **2**: 530-536
- Conete, D., Gava, R., Mestecăneanu, A. (2010) Observations regarding the occurrence of the species *Phalacrocorax pygmeus* Pallas, 1773 and *Aythya nyroca* Linnaeus, 1758 on the basins from the Argeș River (Romania), *Studii și Comunicări, Științele Naturii*, Muzeul Județean Argeș, Pitești, **18**: 105-112
- Conete, D., Mestecăneanu, A., Gava, R. (2006) Speciile de păsări din situl AIA „Lacurile de acumulare de pe Argeș” protejate pe plan național și european, *Argesis, Studii și Comunicări, Științele Naturii*, Muzeul Județean Argeș, Pitești, **14**: 103-115
- Conete, D., Mestecăneanu, A., Gava, R. (2011) The breeding bird species from the middle hydrographical basin of the Arges River (Romania), *Research People and Actual Tasks on Multidisciplinary Sciences*, Lozenec, Bulgaria, **3**: 29-34
- Conete, D., Mestecăneanu, A., Gava, R. (2012) Ornithological researches on the Golești Dam Lake (Argeș County, Romania) during 2003 – 2010, *Analele Universității din Oradea, Fascicula Biologie*, University of Oradea Publishing House, **19** (**1**): 84-92
- Gache, C. (2002) *Dinamica avifaunei în bazinul râului Prut*, Publicațiile Societății Ornitologice Române, Cluj-Napoca, pp. 210
- Gava, R. (1997) Acumulările hidroenergetice de pe râul Argeș, posibile Aree de Importanță Avifaunistică, *Lucrările simpozionului Aree de Importanță Avifaunistică din România*, Publicațiile S. O. R., Cluj-Napoca, **3**: 39-42
- Gava, R., Mestecăneanu, A., Conete, D. (2004a) The reservoirs of the Argeș River valley – important bird areas, *Limnological Reports*, Internat. Assoc. Danube. Res., Novi Sad, Serbia and Montenegro, **35**: 619-631
- Gava, R., Mestecăneanu, A., Conete, D. (2007) The Avifauna of the Middle Basin of Argeș River Artificial Lakes, *Analele Științifice ale Universității „Al. I. Cuza” Iași, s. Biologie animală*, **53**: 187-195
- Gava, R., Mestecăneanu, A., Conete, D. (2008) Avifauna din zona lacurilor de acumulare din bazinul mijlociu și superior al Argeșului și statutul ei de protecție, *Lucrările Conferinței Naționale de Ecologie „Protecția și Restaurarea Bio și Ecodiversității”*, Societatea Română de Ecologie, Ars Docendi, Rezumat extins, pp. 11-13

- Gava, R., Mestecăneanu, A., Conete, D. (2011). Species of birds rarely observed in the Important Bird Area „The Dam lakes of the Argeş River” during of the International waterbird Count (1999 – 2012), *Argesis. Studii și Comunicări, Științele Naturii*, Muzeul Județean Argeş, Pitești, **19**: 79-86
- Gava, R., Mestecăneanu, A., Conete, D. (2012) The mid-winter count of the waterfowls in 2012 on the basins from the Argeş River (IBA – “Lacurile de Acumulare de pe Argeş”). *Scientific Papers. Current Trends in Natural Sciences*, University of Pitești, Faculty of Sciences, **1 (2)**: 6-11
- Gava, R., Mestecăneanu, A., Conete, D., Mestecăneanu, F. (2004b) Recensământul păsărilor de baltă din ianuarie de pe lacurile din bazinul mijlociu al râului Argeş, în perioada 2000 – 2004, *Argesis, Studii și Comunicări, Științele Naturii*, Muzeul Județean Argeş, Pitești, **12**: 125-132
- Mătieș, M. (1969) Cercetări avifenologice de-a lungul bazinului mijlociu și superior al Argeşului între 1 ianuarie – 31 mai 1968, *Studii și Comunicări*, Muzeul Județean Argeş, **2**: 73–90
- Mestecăneanu, A., Gava, R. (2013) The avifauna from Vâlcele, Budeasa, Bascov, Pitești and Golești basins observed in the prevernal season in 2013, *Argesis, Studii și Comunicări, Științele Naturii*, Muzeul Județean Argeş, Pitești, **21**: 71-86
- Mestecăneanu, A., Gava, R. (2015a) The avifauna from Vâlcele, Budeasa, Bascov, Pitești, and Golești dam reservoirs observed in the autumnal season (2013), *Drobeta, Seria Științele Naturii*, Muzeul Regiunii Porților de Fier, Drobeta Turnu Severin, **25**: 103-116
- Mestecăneanu, A., Gava, R. (2015b) The avifauna from Vâlcele, Budeasa, Bascov, Pitești, and Golești dam reservoirs observed in the hiemal season (2013 and 2014), *Oltenia, Studii și comunicări, Științele Naturii*, Muzeul Olteniei Craiova, **31 (1)**: 154-165
- Mestecăneanu, A., Gava, R. (2016a) A year of ornithological observations on the Vâlcele, Budeasa, Bascov, Pitești, and Golești dam reservoirs from ROSPA0062 Lacurile de acumulare de pe Argeş. *Oltenia, Studii și comunicări, Științele Naturii*, Muzeul Olteniei Craiova, **32 (1)**: 97-109
- Mestecăneanu, A., Gava, R. (2016b) The census of the water birds from the dam basins from the Argeş River, between Vâlcele and Golești (January, 2015), *Argesis, Studii și Comunicări, Științele Naturii*, Muzeul Județean Argeş, Pitești, **24**: 81-90
- Mestecăneanu, A., Gava, R. (2016c) The influence of the habitats and anthropogenic pressure on birds, observed during February 2013 – January 2014 on the dam reservoirs from the Argeş River between Vâlcele and Golești, *Scientific Papers, Current Trends in Natural Sciences*, University of Pitești, Faculty of Sciences, **5 (9)**: 18-27
- Mestecăneanu, A., Gava, R. (2017) The Anseriformes from the basins of the Argeş River between Vâlcele and Golești (Argeş County, Romania) in February 2013 – January 2014, *Scientific Papers, Current Trends in Natural Sciences*, University of Pitești, Faculty of Sciences, **6 (11)**: 135-147
- Mestecăneanu, A., Conete, D., Gava, R. (2004) Contribuții la cunoașterea păsărilor clocitoare din bazinul mijlociu al râului Argeş, *Scripta Ornithologica Romaniae*, Cluj Napoca, **1**: 17-20
- Mestecăneanu, A., Conete, D., Gava, R. (2005) Observations of Monitoring Type about the Water Birds from the Golești Accumulation Lake – Argeş River, *Drobeta*, Muzeul Regiunii Porților de Fier, Seria Științele Naturii, Editura Universitaria Craiova, **15**: 114-121

- Mestecăneanu, A., Conete, D., Gava, R. (2006a) The avifauna of Basin Golești, *Acta Universitatis Cibiniensis*, „Lucian Blaga” University of Sibiu, **1, 1 (6)**: 117-126
- Mestecăneanu, A., Conete, D., Gava, R. (2008) Avifauna lacurilor de acumulare de pe râul Argeș dintre Vâlcele și Golești și statutul ei de protecție în baza Directivei Păsări, *ECOS – Revistă de ecologie, educație ecologică și ocrotire a naturii*, Pitești, **20**: 53-61
- Mestecăneanu, A., Conete, D., Gava, R. (2010) Ecological research-studies regarding the avifauna during the hiemal period from the basins area of the Argeș River between 2000 and 2010, *Annals, Food Science and Tehnology*, Universitatea Valahia, Târgoviște, **11, 2**: 127-135
- Mestecăneanu, A., Conete, D., Gava, R. (2013) The midwinter waterbird census from the basins Vâlcele, Budeasa, Bascov, Pitești and Golești from the Argeș River (January 2013), *Scientific Papers, Current Trends in Natural Sciences*. University of Pitești, Faculty of Sciences, **2 (3)**: 51-58
- Mestecăneanu, A., Conete, D., Roșescu, M., Gava, R. (2006b) Observații avifaunistice pe valea râului Argeș, *ECOS – Revistă de ecologie, educație ecologică și ocrotire a naturii*, Pitești, **18**: 68-75
- Mitruly, A. (2002) *Avifauna bazinelor acvatice antropice din Podișul Târnavelor*, Publicațiile Societății Ornitologice Române, Editura Risoprint, Cluj-Napoca, pp. 172
- Munteanu, D. (1978) Avifauna ecosistemelor antropogene și influența presiunii antropice asupra populațiilor de păsări, *Ecosistemele artificiale și însemnătatea lor pentru omenire*, Academia Republicii Socialiste România, Filiala Cluj-Napoca, Subcomisia Om și Biosferă, Cluj-Napoca, **1**: 264-277
- Munteanu, D. (2000) *Avifauna bazinului montan al Bistriței Moldovenești*, Aditura Alma Mater, Cluj-Napoca, pp. 250
- Munteanu, D., Mătieș, M. (1983) Modificări induse de lacurile de acumulare în structura și dinamica avifaunei, *Analele Banatului, Științele Naturii*, Muzeul Banatului, Timișoara, **1**: 217–225
- Munteanu, D., Toniuc, N., Weber, P., Szabó, J., Marinov, M. (1989) Evaluarea efectivelor păsărilor acvatice în cartierele lor de iernare din România (1988, 1989), *Ocrotirea naturii și mediului înconjurător*, București, **33 (2)**: 105–112
- Rang, C. (2002) *Studiul dinamicii unor comunități de păsări din bazinul mijlociu al râului Siret, incluzând zonele lacurilor de acumulare*, Publicațiile Societății Ornitologice Române, Cluj-Napoca, pp. 250
- Truță, A.-M., Dumitru D.-R. (2015) Research on Argeș River fish fauna in Budeasa-Golești Area. *Scientific Papers, Current Trends in Natural Sciences*, Universitatea din Pitești, **4 (8)**: 95-105

Antibiotic susceptibility of bacterial isolates and water quality index of water sourced from closed ground water and open hand dug well in Koko Community, Delta State, Nigeria

Emmanuel Esosa Imarhiagbe¹ and Beckley Ikhajiagbe^{2,✉}

SUMMARY. Water samples were collected from a semi urban community in Nigeria with the aim of investigating the water quality index and antibiotic profile of bacterial isolates of closed ground water and open hand dug wells. Physicochemical and microbiological analyses were carried out using standard analytical methods. pH of groundwater and hand dug well ranged from 4.16 to 5.74 and 4.83 to 5.22 respectively. The total suspended solid of water samples for hand dug well ranged from 1.2-9.2mg/l. Also iron concentrations for groundwater and hand dug well water samples ranged from 0.15-0.54 mg/l and 0.62-1.12 mg/l respectively. Microbial analysis of the water samples revealed the presence of bacteria such as *S. aureus*, *Klebsiella* sp., *E. coli*, *B. subtilis*, *Pseudomonas*, *Aeromonas hydrophila* and *Enterobacter aerogenes* and fungi such as *Aspergillus niger*, *Penicillium notatum*, *Aspergillus flavus*, *Rhizopus stolonifer* and *Saccharomyces cerevisiae*. The total heterotrophic bacterial count of water samples for groundwater and hand dug well samples indicated that values ranged from $2.9 - 4.4 \times 10^3$ cfu/ml and $5.4 - 8.6 \times 10^3$ cfu/ml respectively. Total coliform of water samples for groundwater and hand dug well samples indicated that values ranged from 5-8 MPN/100ml and 10-20 MPN/100ml respectively. *E. coli* count of water samples for groundwater and hand dug well samples indicated that values ranged from 0.0 MPN/100ml and 4-8 MPN/100ml respectively while total fungal count of groundwater and hand dug well samples indicated that values ranged from $0.0 - 6.0 \times 10^2$ cfu/ml and $3.5 \times 10^2 - 17.0 \times 10^3$ cfu/ml respectively. Variable antibiotic susceptibility patterns were observed in antibiotic inhibitory zone (mm) among the tested bacterial isolates. Evaluation of Water Quality Index indicated values of 34.4 for groundwater source indicating good water quality and 67.31 for open hand dug well

¹ Department of Environmental Management and Toxicology, Faculty of Life Sciences, University of Benin, Benin City, Nigeria.

² Environmental Biotechnology and Sustainability Research Group, Department of Plant Biology and Biotechnology, Faculty of Life Sciences, University of Benin, Nigeria.

✉ **Corresponding author: Beckley Ikhajiagbe**, Environmental Biotechnology and Sustainability Research Group, Department of Plant Biology and Biotechnology, University of Benin, Nigeria.
E-mail: beckley.ikhajiagbe@uniben.edu

indicating water was of poor quality. Findings from this study revealed that groundwater sources had better and acceptable quality compared to those of open hand dug wells, hence it is recommended that critical measures be put in place to ensure the safety of both sources of water in Koko community.

Keywords: antibiotic susceptibility, ground water, hand dug wells, microbiological analysis, water quality

Introduction

Research findings have revealed that a considerable percentage of all diseases which cause mortality in the developing countries are directly and indirectly related to poor drinking water quality (Jeffre, 2008), and over 20,000 children die per day (approximately six million annually) due to water borne diseases resulting from availability of safe drinking water (TWAS, 2002). The geological constraints limit accessibility of many human communities to water that is adequate in terms of quantity, quality and sustainability (Tchobanoglous *et al.*, 2003) and lack of adequate supply of potable water is a critical challenge in Nigeria and other developing countries.

In these parts of the world, the usual sources of drinking water are streams, rivers, wells and boreholes which are mostly untreated and associated with various health risks (Agbarie and Obi, 2009). Paucity of infrastructure for effective treatment and distribution of water accounts for the incidence of high morbidity and mortality rate associated with water borne diseases in developing countries (Muhammed *et al.*, 2007). The quality of water influences the health status of any population, hence, analysis of water for physical, biological and chemical properties including trace element contents are very important for public health studies (Orewole *et al.*, 2007).

A reliable supply of clean wholesome water is highly essential in a bid to promote healthy living amongst the inhabitants of any defined geological region (Ndinwa *et al.*, 2012). In advance industrialized world, delivery of safe drinking water and sanitation technologies are, however, not affordable (Ashaye *et al.*, 2001; Adekunle *et al.*, 2004). Consequently, given the renewed global commitment towards the millennium development goals (MDG) marked for 2015, the importance and contribution of locally sourced, low cost alternative drinking water schemes to sustainable access in rural, sub-urban and urban settings of developing countries cannot be over emphasized (UNDESA, 2004).

The objective of this study was to evaluate the antibiotic susceptibility profile of bacterial isolates and water quality index of closed groundwater and open hand dug wells water sourced from Koko community located in Delta State, Nigeria.

Materials and methods

Sample collection

Water samples were collected from closed ground water wells (GW) and open hand dug wells (HDW) at different locations covering the geographic land mass of Koko community, from November to December 2016 and January to March 2017. The water samples were collected using sterile sampling bottles and transported to the laboratory for microbiological and physico-chemical analysis.



Figure 1. Google earth map indicating the respective sampled codified locations in Koko community, Delta State, Nigeria

Determination of heterotrophic bacterial and fungal counts:

Enumeration of the total viable bacterial and fungal counts were determined using serial dilution and pour plate methods as described by Harley and Prescott (2002). Media use was sterile Peptone Water as diluent, Nutrient Agar (for bacterial count) and Potato Dextrose Agar (for fungal count). Plating was done in duplicates and the Nutrient Agar plates were incubated at 35°C for 48h in an incubator while the Potato Dextrose Agar plates were incubated at room temperature for 5 days. Sub-culturing of representatives of the various colonies onto agar plates of the same media were made for purification. The bacterial isolates were Gram-stained (Cheesbrough, 2000). Phenotypic profiling of both Gram-positive and Gram-negative bacteria was

undertaken using API 50CHB and API 20E strips (BioMerieux, Marseille, France). Additional tests of spore stain and oxidase were also performed. Observable colonial characteristics of the fungal isolates which were noted, microscopic observation of portions of their mycelia and spores using wet mount technique were used in identifying the fungal isolates. Observed fungal spores and mycelial fragment were compared to illustrations contained in Barnett and Hunter (1972).

Determination of coliform and *E. coli* counts:

These tests were carried out according to methods stated by Cheesebrough (2006) and were conducted in three stages: presumptive stage, confirmatory stage and completed stage.

Determination of antibiotic susceptibility profile of the bacterial isolates:

The antibiogram of the isolates was determined using the disc diffusion method (Harley and Prescott, 2002). The test bacterial isolates were inoculated unto Muller-Hinton Agar and followed by application of the discs (Oxoid) impregnated with different antibiotics. Antibiotic disc contained the following antibiotics: Ciprofloxacin (CPX, 10 µg), Chloramphenicol (CH, 30 µg), Sparfloxacin (SP, 10 µg), Amoxicillin (AM, 30 µg), Augmentin (AU, 30 µg), Gentamicin (CN, 30 µg), Pefloxacin (PEF, 10 µg), Streptomycin (S, 30 µg), Erythromycin (E, 10 µg), Ampiclox (APX, 30 µg), Zinnacef (Z, 20 µg), Ofloxacin (OFX, 5 µg) and Recephin (R, 25 µg). After 24 hours of incubation at 30 °C, the diameter of the zone of inhibition of each antibiotic disc was measured using a ruler and recorded.

Physicochemical analyses:

pH, electrical conductivity, turbidity, total suspended solids (TSS), total dissolved solids (TDS), dissolved oxygen (DO), biochemical oxygen demand (BOD₅) were analyzed according to standard analytical procedure (APHA, 1993, Ademoroti, 1996).

Evaluation of heavy metals:

The levels of iron (Fe), manganese (Mn), zinc (Zn), copper (Cu), chromium (Cr) and lead (Pb) were analysed using the atomic absorption spectrophotometer (AAS) (Buck Scientific, Model 210 VGP).

Evaluation of the water quality index of the sampled waters:

The WQI scores were generated using the Weighted Arithmetic Index method as described by Chatterjee and Raziuddin (2002).

$$WQI = \{[(V_{\text{actual}} - V_{\text{ideal}}) / (V_{\text{standard}} - V_{\text{ideal}})] \times 100\}$$

Table 1.

Water Quality Index (WQI) rankings

Range	Ranking/Status
0-25	Excellent
26-50	Good
51-75	Poor
76-100	Very poor
Above 100	Unsuitable for drinking

Source: Chatterjee and Raziuddin (2002)**Statistical analysis**

The non-parametric analogue of the unpaired student T test was utilized to determine if the differences between the respective microbial and physicochemical data recorded for the ground water and hand dug well water samples was statistically significant ($p < 0.05$).

The results of mean heterotrophic bacterial and fungal counts were in the order 10^2 to 10^3 cfu/ml respectively. Generally, water samples from open hand dug wells had higher counts (5.4×10^3 to 8.6×10^3 cfu/ml) than ground water (table 2). Total microbial counts are important parameters which for indicative of the hygienic and portability properties of drinking water.

Table 2.

The heterotrophic microbial and coliform counts of the water samples

Sampling month	Water type	Mean HBC ($\times 10^3$ cfu/ml)	Mean HFC ($\times 10^2$ cfu/ml)	TCC (MPN/100ml)	<i>E. coli</i> count (MPN/100ml)
November 2016	GW	4.4	0.0	5	0.01
	HDW	6.3	5.5	10	5
December 2016	GW	3.6	0.0	5	0.0
	HDW	5.4	3.5	10	4
January 2017	GW	4.2	4.0	7	0.0
	HDW	6.5	7.5	10	5
February 2017	GW	2.9	4.0	5	0.0
	HDW	8.0	9.0	10	4
March 2017	GW	3.7	6.0	8	0.0
	HDW	8.6	17.0	20	8
p-values	-	0.022	0.000	0.015	0.000

KEY: GW = Closed groundwater sample, HDW = hand dug well, HBC = Heterotrophic bacterial count, HFC = Heterotrophic fungal count, TCC = Total coliform count

The results showed that the greatest frequency was observed for *Bacillus* spp. (30.1 %) and the least predominant among the bacterial isolates was *Escherichia coli* (6.0 %). Regarding the fungal isolates *Aspergillus flavus* was the most predominant (26.5 %) and *Rhizopus stolonifer* (11.8 %) was the least predominant (table 3). Bacterial isolates identified in the water samples include: *Micrococcus* sp., *Klebsiella* sp., *Escherichia coli*, *Bacillus* sp., *Pseudomonas aeruginosa* and *Aeromonas hydrophila*. Fungal isolates identified in the water samples collected include *Aspergillus niger*, *Penicillium notatum*, *Aspergillus flavus*, *Rhizopus stolonifer* and *Saccharomyces cerevisiae*. These microbial organisms are of public health importance. The high microbial load and variety of microorganisms detected and isolated from these water samples may be attributed to the poor sanitary and water handling practices.

Table 3.

Percentage frequency of occurrence of bacterial and fungal isolates from the respective water samples

Bacterial isolate	No. of bacterial isolates	% frequency of occurrence	Fungal isolate	No. of fungal isolates	% frequency of occurrence
<i>Micrococcus</i> spp.	18	21.7	<i>Aspergillus niger</i>	8	23.5
<i>Klebsiella</i> spp.	7	8.4	<i>Penicillium notatum</i>	8	23.5
<i>Escherichia coli</i>	5	6.0	<i>Aspergillus flavus</i>	9	26.5
<i>Bacillus</i> spp.	25	30.1	<i>Rhizopus stolonifera</i>	4	11.8
<i>Pseudomonas aeruginosa</i>	8	9.6	<i>Saccharomyces cerevisiae</i>	5	14.7
<i>Aeromonas hydrophila</i>	20	24.1			
	n =83	100		n =34	100

Variable antibiotic susceptibility patterns were observed in the antibiotic inhibitory zones (mm) among the tested bacterial isolates (Table 4). Findings showed that seven (7) antibiotics did not exhibit inhibition zones against tested Gram negative bacterial isolates (*Klebsiella* sp., *Aeromonas hydrophila*, *E.coli* and *Pseudomonas aeruginosa*) where as five (5) antibiotics showed no inhibition against Gram positive bacterial isolates (*Micrococcus* and *Bacillus*). The widest zone of inhibition was observed for ofloxacin (30.0 mm) against *Klebsiella* sp., which indicates susceptibility of the bacterial isolate. This observation could be ascribed to the fact that many bacteria isolated from the surface water possess an important ecological quality, namely that of resistance plasmids, which can be picked up in the course of selective processes. Researches had reported the increased incidences of bacterial resistance to antibiotics which contribute to the prevailing trend of antibiotic abuse and misuse by a larger human population most especially in the developing world (Omogbai and Ikenebomeh, 2013, Imarhiagbe *et al.*, 2016). Spanggard *et al.* (1993) had earlier observed the possibility of members of bacterial groups residing in a common environment to express a similar antibiotics pattern if they share in a common pool of R-factor plasmids.

Table 4.

Zone(s) of inhibition (mm) representing antibiotic sensitivity/resistance patterns elicited by the bacterial isolates against the test drugs

Gram +ve	PEF (10 µg)	CN (30 µg)	APX (30 µg)	Z (20 µg)	AM (30 µg)	R (25 µg)	CPX (10 µg)	S (30 µg)	SXT (30 µg)	E (10 µg)
(1)	0.0	13.0	0.0	17.0	0.0	20.0	24.0	0.0	0.0	11.0
(2)	19.0	25.0	0.0	20.0	12.0	28.0	30.0	14.0	0.0	18.0
Gram -ve	SXT(30 µg)	CH (30 µg)	SP (10 µg)	CPX (10 µg)	AM (30 µg)	AU (30 µg)	CN (30 µg)	PEF (10 µg)	OFX (5 µg)	S (30 µg)
(3)	0.0	08.0	0.0	18.0	0.0	14.0	20.0	0.0	27.0	15.0
(4)	0.0	0.0	0.0	15.0	0.0	0.0	16.0	0.0	20.0	0.0
(5)	0.0	0.0	0.0	20.0	0.0	0.0	18.0	0.0	25.0	0.0
(6)	0.0	13.0	15.0	22.0	0.0	12.0	20.0	10.0	30.0	10.0

KEY: (1) *Micrococcus* sp., (2) *Bacillus* sp., (3) *Aeromonas hydrophila*, (4) *Pseudomonas aeruginosa*, (5) *E. coli*, (6) *Klebsiella* sp. Ciprofloxacin (CPX), Chloramphenicol (CH), Sparfloxacin (SP), Amoxicillin (AM), Augmentin (AU), Gentamicin (CN), Pefloxacin (PEF), Streptomycin (S), Erythromycin (E), Ampiclox (APX), Zinnacef (Z), Ofloxacin (OFX) and Recephin (R) S= Streptomycin.

Table 5 revealed the result of the physicochemical analysis of the water samples from closed groundwater and open hand dug well from Koko community. pH value is significant determinant for the suitability of water for several purposes. pH of the groundwater samples and hand dug well water samples ranged from 4.40-5.741 and 4.83-5.54 respectively. The findings show that water from this location is slightly acidic; this observation may be due to the geological conditions (King and Ekeh, 1990). Electrical conductivity for groundwater samples and hand dug well water samples ranged from 52.7-86.2 µS/cm and 145.3-330 µS/cm respectively. Measurement of turbidity has been described as a very important parameter when evaluating the quality of a drinking water. Findings from this study revealed that water sourced from closed ground water and open hand dug wells had a mean 0.0 and 2.82 NTU respectively. The highest values of TDS and TSS were observed in open hand dug wells (4.54 mg/L and 98.10 mg/L respectively). Dissolved oxygen for both groundwater samples and hand dug well water samples ranged from 5.5-7.1 mg/L and 4.5-6.6 mg/L respectively. BOD₅ for groundwater samples and hand dug well water samples ranged from 0.4-1.1mg/L and 1.9-4.7mg/L. Also heavy metal analysis of water samples from closed groundwater and open hand dug wells is indicated in Table 5.

Table 5.

Physicochemical analysis of water samples from closed Groundwater (GW) and open Hand Dug (HDW) wells from Koko community

Parameter	Water type	Month					(F)	(G)
		(A)	(B)	(C)	(D)	(E)		
pH	GW	5.19	5.74	4.54	4.40	4.16	4.81±0.65	6.5-8.5
	HDW	5.54	4.83	5.12	4.92	5.22	5.13±0.28	
EC (µS/cm)	GW	60	57.8	63.8	52.7	86.2	64.1±12.99	NA
	HDW	330	152.6	174.1	145.3	193.1	199.02±75.6	
Turbidity (NTU)	GW	0.0	0.0	0.0	0.0	0.0	0.00±0.00	5.0
	HDW	0.0	2.7	8.8	0.0	2.6	2.82±3.59	
TSS (mg/l)	GW	0.0	0.0	0.0	0.0	0.0	0.00±0.00	500
	HDW	1.2	5.1	9.2	2.1	5.1	4.54±3.14	
TDS (mg/l)	GW	31	28.9	31.7	28.1	42.9	32.52±5.99	500
	HDW	160	76.1	86.3	72.3	95.8	98.10±35.79	
DO (mg/l)	GW	5.9	6.3	5.5	6.8	7.1	6.32±0.65	NA
	HDW	6.1	4.5	5.2	5.9	6.6	5.66±0.82	
BOD ₅ (mg/l)	GW	0.4	0.0	0.0	1.5	1.1	0.60±0.67	NA
	HDW	4.7	2.3	1.9	3.7	4.8	3.48±1.34	
p-values	-	0.000	0.000	0.000	0.000	0.000	0.000	-

(A) November 2016, (B) December, 2016, (C) January, 2017, (D) February 2017, (E) March 2017, (F) Mean ± S.D, (G) WHO limit, (NA) Not available

Table 6.

Water Quality Index of closed groundwater and open hand dug sources

Parameter	Closed ground water (GW)					Open hand dug wells (HDW)				
	V _i	S _i	W _i	q _i	W _i q _i	V _i	S _i	W _i	q _i	W _i q _i
pH	4.81	6.5-8.5	0.22	0	0	5.13	6.5-8.5	0.22	0	0
EC (µS/cm)	64.1	250	0.37	25.6	9.51	199.02	250	0.37	79.61	29.54
TDS (mg/L)	32.5	500	0.004	6.50	0.02	98.10	500	0.004	19.62	0.07
TSS (mg/L)	0.00	500	0.003	0	0	4.54	500	0.004	0.91	0.003
DO (mg/L)	6.32	5	0.37	86.3	32.11	5.66	5	0.37	93.13	34.67
BOD ₅ (mg/L)	0.60	5	0.37	12.0	4.47	3.48	5	0.37	69.60	25.91
			ΣW _i =		ΣW _i q _i =			ΣW _i =		ΣW _i q _i =
			1.34		46.1			1.34		90.2
WQI	$(\Sigma W_i q_i / \Sigma W_i) = 46.1/1.34 = 34.4$					$(\Sigma W_i q_i / \Sigma W_i) = 90.2/1.34 = 67.31$				
Ranking	Good					Poor				

Observed (V_i), Standard values (S_i), Unit weights (q_i), and quality rating (q_i)

The observed water quality was supported by the findings from the analytical physico-chemical parameters (Table 6). Water quality index (WQI) value obtained closed groundwater sources and open hand dug wells was 34.4 and 67.31 respectively. The derived water quality index of the respective water

samples revealed that the overall water quality of the samples sourced from open hand dug wells was poor as compared with closed ground water. Based on the WQI values of the samples, it can be inferred that the water samples from open hand dug wells are unfit for direct consumption by the inhabitants of this community.

Table 7.

Levels of heavy metals (mg/l) in water samples collected from closed groundwater and open hand dug wells sited in Koko town, Delta State

Parameter	Water type	Month					(F)	(G)
		(A)	(B)	(C)	(D)	(E)		
Fe	GW	0.28	0.15	0.54	0.26	0.17	0.28±0.156	0.30
	HDW	0.62	0.79	1.12	0.84	0.62	0.80±0.205	
Mn	GW	0.01	0.01	ND	0.01	ND	0.004±0.005	0.1
	HDW	0.05	0.01	0.14	0.09	0.055	0.07±0.049	
Zn	GW	0.08	0.03	0.18	0.07	0.09	0.09±0.055	5.00
	HDW	0.17	0.24	0.62	0.45	0.13	0.32±0.207	
Cu	GW	ND	0.01	ND	0.004	ND	0.0028±0.004	1.0
	HDW	0.02	0.02	0.04	0.061	0.027	0.03±0.017	
Cr	GW	ND	ND	ND	ND	ND	0.00±0.00	0.050
	HDW	0.003	ND	0.05	0.032	0.019	0.02±0.021	
Pb	GW	ND	ND	ND	ND	ND	0.00±0.00	0.01
	HDW	ND	ND	0.02	0.005	ND	0.005±0.009	
p-values	-	0.001	0.001	0.001	0.005	0.001	0.001	

Key: (A) November 2016, (B) December, 2016, (C) January, 2017, (D) February 2017, (E) March 2017, (F) Mean ± S.D, (G) WHO limit, iron (Fe), manganese (Mn), zinc (Zn), copper (Cu), chromium (Cr) and lead (Pb)

Generally, heavy metal levels were below recommended permissible levels by World Health Organisation for drinking water. Iron concentration in both groundwater samples and hand dug well water samples ranged from 0.15-0.28mg/L and 0.62-1.12 mg/L (Table 7).

Conclusions

The results obtained from this research indicate that the water samples collected from this community had poor microbiological and overall Water Quality indices and therefore is unsuitable for direct human consumption. Several measures which include boiling, filtration and addition of flocculants such as aluminium salts would invariably enhance the potability of the water and reduce the risk of developing

water borne gastroenteritis which could arise from the direct consumption of the contaminated water. The entitled authorities ought to ensure proper awareness for treatment of the water sources to safeguard the public health of the population.

REFERENCES

- Adekunle, L. V., Sridhar, M. K., Ajayi, A. A., Oluwale, P. A., Olawuyi, J. F. (2004) An assessment of health and socio-economic implications of sachet in Ibadan, Nigeria: A Public Health Challenge. *African Journal of Biomedical Research* **7**(1): 5–8
- Ademoroti, C. M. A. (1996) *Standard Methods for Water and Effluent Analysis*. Foludex Press. Ltd. Ibadan, 182 pp.
- Agbaire, P. O., Obi, C. G. (2009) Seasonal variations of some physico-chemical properties of River Ethiopie Water in Abraka, Nigeria. *Journal of Applied Science and Environmental Management* **13**(1): 55-57
- American Public Health Authority (APHA) (1993) *Standard Methods for Examination of Water and Wastewater*. 18th Edn. APHA. Washington D.C. 1421 pp.
- Ashaye, O. A., Couple, A. A., Afolabi, O. O., Fasoyiro, S. B. (2001) Physicochemical properties of pure water samples in South Western, Nigeria. *Journal of Food Technology* **6**(4): 119–120
- Barnett, H. L., Hunter, B. B. (1972) *Illustrated Genera of Imperfect Fungi*. 3rd edn. Burgess Publishing Co. New York, 225 pp.
- Chatterjee, C., Raziuddin, M. (2002) Determination of water quality index of a degraded river in Asanol industrial area, Raniganj, Burdwan, West Bengal. *Nature, Environment and Pollution Technology* **1**(2): 181-189
- Cheesebrough, M. (2006) *District Laboratory Practice in Tropical Countries*. Cambridge University Press. London, 434 pp.
- Harley, J.P. and Prescott, L.M. (2002). *Laboratory Exercises in Microbiology*. 5th Edn. Mac Graw Hill, New York, 449 pp.
- Imarhiagbe, E. E, Obayagbona, O. N, Osarenotor, O., Eghomwanre, A. F. (2016) Antibiotic sensitivity pattern of bacterial isolates and physico-chemical composition of maize flour sold in major markets in Benin City, Midwestern Nigeria, *Studia UBB Biologia*, **61**(2): 5-12
- Jeffre, H. (2008) Water Problems, Solutions and Conservation in the Developing World <http://factsanddetails.com>
- King, R. P., Ekeh, I. B. (1990) The status and seasonality in the physico-chemical hydrology of a Nigerian head water stream. *Acta Hydrobiol.*, **32**, 313-328
- Muhammad, B. G., Ismail, B. S., Ekhwan, T., Sujaul, I. M., Tan, C. C. (2007) A physicochemical assessment of the Behar River, Pahang, Malaysia. *Global Journal of Environmental Research* **1**: 7-11

- Nwodo, C. S., Obinna, C. N., Adetayo, Y., Oluwadamisi, V. N. (2011) Assessment of water quality in Canaanland, Ota, Southwest Nigeria. *Agriculture and Biology Journal* **2**(4): 577-583
- Omogbai, B. A., Ikenebomeh, M. (2013) Microbiological characteristics and phytochemical screening of some herbal teas in Nigeria, *Euro. Sci. J.* **18**, 149-160
- Orewole, M. O., Mkaide, O. W., Adekalu, K., Shittu K. A. (2007) Chemical examination of piped water supply of Ile-Ife in South West Nigeria. *Iran Journal of Environment Health Science* **4**(1): 51-56
- Spanggard, B., Jorgenses, F. G., Huss, H. H. (1993) Antibiotic resistance in bacteria isolated from three freshwater farms and an unpolluted stream in Denmark, *Aquaculture* 195-207
- Tchobanoglous, G., Burton, F. L., Stensel, H. D. (2003) *Wastewater Engineering (Treatment Disposal Reuse) / Metcalf & Eddy, Inc.* (4th Edition ed.). McGraw-Hill Book Company, 88pp.
- Third World Academy of Sciences (TWAS) (2002) Safe Drinking Water: the need, the problem, solutions and action plan, Trieste, Italy. <http://www.g77.org/sshlest/TWAS>
- UN Department of Economic and Social Affairs (UNDESA) (2004) Urban agglomerations. Population Division of the Department of Economic and Social Affairs, United Nations, 78pp.
- WHO (2006) Guidelines for drinking Water Quality: Recommendations. 3rd Edition, WHO. Geneva, 66pp.

Preliminary data regarding genetic diversity of several endangered and endemic *Dianthus* species from Romania generated by RAPD markers

Anca-Livia Butiuc-Keul^{1,2,✉}, Liliana Jarda³, Irina Goia^{2,4}, Irina Holobiuc⁵,
Anca Farkas^{1,2} and Victoria Cristea^{3,2}

SUMMARY. Conservation of endangered and endemic species of *Dianthus* from Romania, requires the investigation of genetic polymorphism in the populations. Preliminary data were obtained by molecular characterization using RAPD markers. DNA amplification with the 9 RAPD primers of the individuals belonging to different populations of *D. callizonus*, *D. giganteus* ssp. *banaticus*, *D. glacialis* ssp. *gelidus*, *D. henteri*, *D. nardiformis*, *D. pratensis* ssp. *racovitzae*, *D. spiculifolius* and *D. tenuifolius* revealed low level of polymorphism within and between populations. Several polymorphic RAPD markers were identified being useful for investigation of genetic diversity. Out the 9 primers studied by us, only the primer OPB-07 ensured amplification in all species and primers OPA-13, OPE-04 and 1225 showed positive results in most of the species. The primers 4A-26 and 4A-27 ensured amplification only in *D. spiculifolius* and the primers 4A-23 and OPM-18 gave no results in none of the species.

Keywords: *Dianthus*, endemic/endangered species, genetic polymorphism, RAPD markers.

Introduction

Nowadays, in response to an alarming increase of species number with different degrees of endangerment, the conservation of biodiversity has a great

¹ Babeș-Bolyai University, Faculty of Biology and Geology, Department of Molecular Biology and Biotechnology, 400084 Cluj-Napoca, Romania.

² Babeș-Bolyai University, Center of Systems Biology, Biodiversity and Bioresources, Laboratory of Plant Biology, Genetics, Microbiology and Biotechnology, 400006 Cluj-Napoca, Romania.

³ Babeș-Bolyai University, Alexandru Borza Botanical Garden, 400015 Cluj-Napoca, Romania.

⁴ Babeș-Bolyai University, Faculty of Biology and Geology, Department of Taxonomy and Ecology, 400015 Cluj-Napoca, Romania.

⁵ Institute of Biology, Romanian Academy, 060031 Bucharest, Romania.

✉ **Corresponding author:** Anca-Livia Butiuc-Keul, Babeș-Bolyai University, Faculty of Biology and Geology, Department of Molecular Biology and Biotechnology, 400084 Cluj-Napoca, Romania, E-mail: anca.keul@ubbcluj.ro

importance. The concept of sustainable development implies the conservation of biodiversity both in situ and ex situ. Ex situ conservation is a viable alternative for highly endangered taxa in their natural environment, or may complete the in situ conservation methods (Marriott and Sarasan, 2010; Johnson *et al.*, 2012; Butiuc-Keul, 2006; 2014; Cristea *et al.*, 2014) that has consequences not only in science, but also in economy and culture. In order to develop a proper conservation management it is very important to know the genetic variability of in situ populations. In the case of in vitro conservation, it is also necessary to evaluate the somaclonal variability (i.e., the variability induced by the in vitro conditions) and to avoid the genetic uniformity. According to the International Union for Conservation of Nature (IUCN) there are approximately 33400 plant species threatened with extinction (Primack *et al.*, 2008). Moreover, in 2014, there were 18291 plant species on IUCN red list (<http://discover.iucnredlist.org/search?key=plants>).

Genetic structure of plant populations, the level of genetic polymorphism within and among populations could offer valuable information in order to develop proper strategies for their conservation. The aim of conservation programs is to preserve a high number of populations, as well as their genetic structure and variability (Halmagyi and Butiuc-Keul, 2007; Butiuc-Keul, 2014). A characteristic of endangered and/or endemic plants is the maintenance of low levels of genetic variation in populations. Limited genetic diversity has been reported for many endangered and endemic plant species. The genetic structure and variability of endemic and rare plant populations were intensively studied by DNA markers (Luan *et al.*, 2006; Breinholt *et al.*, 2007; Cristea *et al.*, 2014; Jarda *et al.*, 2014).

Dianthus is one of the most diverse Mediterranean genera, more than 300 species were identified in Eurasia and Africa and over 100 species were found in Europe, more than 70 being endemic (Valente *et al.*, 2010). In Romania, 58 *Dianthus* taxa have been recorded (Ciocârlan, 2009), of which 8 are endemic.

Our previous studies regarding the conservation of endemic and endangered species ensured valuable data for ex situ conservation strategies of several plant species of *Dianthus* as *D. giganteus* ssp. *banaticus* (Jarda *et al.*, 2014), *D. henteri* (Cristea *et al.*, 2010), *D. spiculifolius* (Butiuc-Keul *et al.*, 2001; 2016; Cristea *et al.*, 2006; 2009; 2013; 2014). Concerning the *Dianthus* genus our major interest was focused on the evaluation of genetic structure of populations and genetic variability within and between populations of several endemic and endangered species from Romania. Molecular markers as RAPD are valuable tools for such investigations because they are cost efficient, requires lower amounts of DNA and there is no necessary information about marker sequence. Thus, in this study we report the preliminary data regarding the usefulness of RAPD markers for the assessment of genetic polymorphism in the population of several *Dianthus* species from Romania.

Materials and methods

Plant material

The plant material was collected from different populations of the studied *Dianthus* taxa from Romania. Five individuals were collected from each location. Thus, *D. callizonus* was collected from 2 locations from Piatra Craiului Mountains: P1-Spîrlea refuge and Zăplaz (45°31'42.604"N, 25°12'00.174"E) and Diana refuge and Padina Popii, Piatra Craiului Mountains (45°33'29.184"N, 25°14'29.754"E); *D. giganteus* ssp. *banaticus* was collected from 3 populations: P1-Eșelnița (44°39'27.99"N, 22°17'47.42"E), P2-Domogled Mt. (44°52'36.20"N, 22°26'13.00"E) and P3-Țesna Gorge (44°58'03.30"N, 22°30'34.10"E); *D. glacialis* ssp. *gelidus* from 3 populations from Bucegi Mountains: P1-Omu Peak (45°22'51.004"N, 25°30'40.004"E), P2-Bâlea Lake, Făgăraș Mountains (45°31'40.004"N, 24°44'26.004"E), P3-Obârșia, Bucegi Mountains (45°22'50.004"N, 25°30'39.004"E); *D. henteri* from 3 populations from Vâlcea county: P1-Cornet (45°23'19.82"N, 24°18'28.54"E), P2-Călinești Valey (45°22'19.34"N, 24°17'23.23"E), P3-Jiului Gorge (45°16'46.004"N, 23°23'19.004"E); *D. nardiformis* from 3 populations from Tulcea county: P1-Allah Bair (44°29'01.42"N, 28°13'24.88"E), P2-Consul Hill (45°10'55.19"N; 28°16'15.77"E), P3-Măcin (45°14'22.71"N, 28°35'01.01"E); *D. pratensis* ssp. *racovitzae* from 2 populations: P1-Dorobanț village, Iassy County (47°14'33"N, 27°34'55"E), P2-Spătaru, Buzău County (45°4'14.56"N, 26°47'1.64"E); *D. spiculifolius* from 2 populations: P1-ROSCI0027 Natura 2000 site, Hășmaș Mountains (46°44'274"N, 25°47'584"E), P2-ROSCI0002 Natura 2000 site, Apuseni Natural Park, Vlădeasa Massif (N 46°35'454"N, 22°48'384"E); *D. tenuifolius* from 2 populations from Suceava county: P1-Stânișoarei Mountains (47°21'50.87"N, 25°36'10.88"E), P2-Bistriței Mountains (47°23'30.00"N, 25°30'16.65"E).

RAPD analysis

DNA was isolated using the CTAB method (Doyle and Doyle, 1987), RAPD markers were obtained by PCR amplification, performed in 25 μ L of mixture containing 2 mM MgCl₂, 1 μ M of each primer, 200 μ M of each dNTP, 1.0 U of Taq (Fermentas) in reaction buffer (10mM TrisHCl pH 8.8, 50 mM KCl, 1.5 mM MgCl₂) and 25 ng of genomic DNA. Amplification programme: 1. T=94°C, 4 min; 2. T=94°C, 45 s; 3. primer alignment at 36 °C, 45 s; 4. elongation T=72°C, 50 s; steps 2-4, 35X. 9 primers have been tested as it can be seen in Table 1. Amplicons were separated on 1.5% agarose gel, stained with 0.5 μ g mL⁻¹ ethidium bromide. At least 2 independent PCR amplifications were performed for each primer.

Genetic similarities between individuals and populations were measured by Euclidian distance with the Past programme and the generated similarity coefficients were used for constructing a dendrogram with UPGMA option using the same programme.

Table 1.

Characteristics of primers used for RAPD amplification

No.	Primer sequence	Temperature of alignment
1	1225: 5'-AGGTGACCGT-3'	36 °C
2	OPA-13: 5'-CAGCACCCAC-3'	36 °C
3	OPB-07: 5'-GGTGACGCAG-3'	36 °C
4	OPE-04: 5'-GTGACATGCC-3'	36 °C
5	OPF-04: 5'-GGTGATCAGG-3'	36 °C
6	OPM-18: 5'-CACCATCCGT-3'	36 °C
7	4A-23: 5'-TCGCGAGCTG-3'	36 °C
8	4A-26: 5'-GTGATCGCAG-3'	36 °C
9	4A-27: 5'-CAATCGCCGT-3'	36 °C

Results and discussion

Amplification with the RAPD primers allows the identification of several polymorphic markers useful for investigation of genetic diversity between and within population. Out of the 9 primers studied by us, only the primer OPB-07 ensured amplification in all species. The specific patterns of amplification with this primer in *Dianthus* species are shown in Fig. 1.

The primer OPA-13 showed results in all species except *D. nardiformis*. The primers 1225 ensured amplification in most of the species except *D. henteri* and *D. pratensis* ssp. *racovitzae* and the primer OPE-04 also ensured amplification in most of the species except *D. giganteus* ssp. *gelidus* and *D. nardiformis*. Using the primer OPF-04 we obtained the successful amplification only in *D. callizonus*, *D. giganteus* ssp. *banaticus*, *D. giganteus* ssp. *gelidus*, *D. spiculifolius* and *D. tenuifolius*. The primers 4A-26 and 4A-27 showed amplification only in *D. spiculifolius* and the primers 4A-23 and OPM-18 gave no results in none of the species (Table 2). Regarding the polymorphism within and between populations, the results are different in the 8 studied *Dianthus* species.




Figure 1. RAPD patterns of different *Dianthus* species with OPB-07 primer (a- *D.callizonus*: 1-molecular marker, 2-Dc 1.1, 3-Dc 1.2, 4-Dc 1.3, 5-Dc 1.4, 6-Dc 1.5, 7-Dc 1.6, 8-Dc 1.7, 9-Dc 1.8; b-*Dianthus giganteus* ssp. *banaticus*: 1-molecular marker, 2-Dgb 1.1, 3-Dgb 1.2, 4-Dgb 1.3, 5-Dgb 2.1, 6-Dgb 2.2, 7-Dgb 2.3, 8-Dgb 3.1, 9-Dgb 3.2, 10-Dgb 3.3; c- *Dianthus giganteus* ssp. *gelidus*: 1-molecular marker, 2-Dgg 1.1, 3-Dgg 1.2, 4-Dgg 1.3, 5-Dgg 1.4, 6-Dgg 1.5, 7-Dgg 2.1, 8-Dgg 2.2, 9-Dgg 2.3, 10-Dgg 2.4, 11-Dgg 2.5, 12-Dgg 3.1, 13-Dgg 3.2, 14-Dgg 3.3, 15-Dgg 3.4, 16-Dgg 3.5; d- *Dianthus henteri*: 1-molecular marker, 2-Dh 1.1, 3-Dh 1.2, 4-Dh 1.3, 5-Dh 1.4, 6-Dh 1.5, 7-Dh 2.1, 8-Dh 2.2, 9-Dh 2.3, 10-Dh 2.4, 11-Dh 2.5, 12-Dh 3.1, 13-Dh 3.2, 14-Dh 3.3, 15-Dh 3.4, 16-Dh 3.5; e- *Dianthus nardiformis*: 1-molecular marker, 2-Dn 1.1, 3-Dn 1.2, 4-Dn 1.3, 5-Dn 1.4, 6-Dn 1.6, 7-Dn 2.1, 8-Dn 2.2, 9-Dn 2.3, 10-Dn 3.1, 11-Dn 3.2, 12-Dn 3.3; f-*Dianthus pratensis* ssp. *racovitzae*: 1-molecular marker, 2-Dpr 1.1, 3-Dpr 1.2, 4-Dpr 1.3, 5-Dpr 1.4, 6-Dpr 1.6, 7-Dpr 2.1; g-*Dianthus spiculifolius*: 1-molecular marker, 2-6-negative controls, 7-Ds 1.1, 8-Ds 1.2, 9-Ds 1.3, 10-Ds 1.4, 11-Ds 1.5, 12-Ds 2.1, 13-Ds 2.2, 14-Ds 2.3, 15-Ds 2.4, 16-Ds 2.5; h- *Dianthus tenuifolius*: 1-molecular marker, 2-Dt 1.1, 3-Dt 1.2, 4-Dt 1.3, 5-Dt 2.1, 6-Dt 2.2, 7-Dt 2.3; number significance: first number-population, second number-individual). Separation on 1.5% agarose gel stained with 0.5 µg/mL ethidium bromide.

Genetic variability in population of *D. callizonus* was assessed by RAPD markers generated with different primers as 1225, OPE-04, OPF-04, OPB-07 and OPA-13 (Table 2). All primers generated different polymorphic patterns but the highest polymorphism was observed with the primers OPE-04 and OPB-07. The dendrogram constructed based on Euclidian distance between individuals showed that they are clustered in two groups of similarity (Fig. 2).

Table 2.

Genetic polymorphism of *Dianthus* species based on RAPD markers

Species	Population	RAPD fragments						
		1225	OPA-13	OPB-07	OPE-04	OPF-04	4A-26	4A-27
<i>D. callizonus</i>	P1	5	12	7	5	3	-	-
<i>D. giganteus</i> ssp. <i>banaticus</i>	P1	3	5	3	2	5	-	-
	P2	2	7	3	2	3	-	-
	P3	3	4	3	2	3	-	-
<i>D. giganteus</i> ssp. <i>gelidus</i>	P1	4	3	3	-	4	-	-
	P2	2	3	4	-	6	-	-
	P3	4	3	4	-	5	-	-
<i>D. henteri</i>	P1	-	3	2	5	-	-	-
	P2	-	4	2	4	-	-	-
	P3	-	4	4	4	-	-	-
<i>D. nardiformis</i>	P1	1	-	1	-	-	-	-
	P2	2	-	1	-	-	-	-
	P3	3	-	1	-	-	-	-
<i>D. pratensis</i> ssp. <i>racovitzae</i>	P1	-	1	4	3	-	-	-
	P2	-	1	4	1	-	-	-
<i>D. spiculifolius</i>	P1	5	6	5	5	4	2	4
	P2	2	4	2	5	3	2	3
<i>D. tenuifolius</i>	P1	2	3	1	2	2	-	-
	P2	2	3	2	3	5	-	-




Figure 2. Dendrogram illustrating similarities between the individuals of *Dianthus callizonus* based on RAPD markers.

In *Dianthus giganteus* ssp. *banaticus*, RAPD amplification was successfully obtained only with the primers 1225, OPE-04, OPF-04, OPB-07 and OPA-13 (Table 2). The primer 1225 showed 3 bands in the populations P1 and P3, and only 2 bands in the population P2. There is one specific band which is present only in population P3-Cheile Țesnei, Mehedinți. The populations from Eșelnița and Domogled are extremely similar, the individuals showing the same RAPD pattern with this primer. Similar RAPD patterns were also obtained in all populations with the primer OPE-04. The amplification with the primer OPF-04 showed differences between individuals and populations, thus this primers is appropriate for genetic polymorphism assessment. RAPD markers generated with the primer OPB-07 are present in all individuals from P2 population and in some individuals from P1 and P3 populations, these patterns being low polymorphic. High genetic polymorphism was showed by OPA-13 primer, while higher polymorphism being identified in the populations P1 and P2 than in the population P3. Based on RAPD markers a dendrogram of similarity between individuals was generated (Fig. 3). Thus, the individuals of *Dianthus giganteus* ssp. *banaticus* are clustering in 3 groups. All groups are including individuals from all populations. The populations P1 and P3 are more similar than the population P2. In conclusion the primers OPF-04 and OPA-13 showed high genetic polymorphism between and within populations. The primer OPB-07 showed only genetic polymorphism within populations.




Figure 3. Dendrogram illustrating similarities between the individuals of *Dianthus giganteus* ssp. *banaticus* based on RAPD markers.

RAPD amplification of *Dianthus giganteus* ssp *gelidus* was successfully obtained with primers 1225, OPF-04, OPB-07 and OPA-13 (Table 2). Amplification with 1225 primer generated 7 different bands showing high level of genetic polymorphism between and within populations. The OPF-04 primer generated high number of bands with polymorphic distribution between populations. The OPB-07 primer generated 5 different bands that showed genetic polymorphism between populations. The primer OPA-13 showed the same pattern of the individuals from P3 population and low genetic polymorphism between individuals from P1 and P2 populations. The dendrogram of similarity showed that the individuals from P3 population and several individuals from P1 population are clustered together, while the individuals from P2 population are clustered together in a distinct group. Several individuals from P1 population are clustered in other group at higher genetic distance (Fig. 4).

RAPD amplification in *D. henteri* was successfully obtained only with the following primers OPE-04, OPB-07 and OPA-13 (Table 2). DNA amplification with the primer OPE-04 generated 5 bands that are present in several individuals, thus this primer showed low genetic polymorphism between and within populations. Similar results were also been obtained with the primer OPB-07, 4 different bands being obtained that are present in some of the individuals. High genetic polymorphism was observed by DNA amplification with the primer OPA-13, 4 fragments were observed but their distribution is very different in the individuals belonging to this plant species. Similarity dendrogram constructed based on RAPD markers showed that the individuals from all populations are very similar. The individuals are clustered in 3 groups, the first and the second contain most of the individuals, while the third group contains only few individuals (Fig. 5).




Figure 4. Dendrogram illustrating similarities between the individuals of *Dianthus giganteus* ssp. *gelidus* based on RAPD markers.




Figure 5. Dendrogram illustrating similarities between the individuals of *Dianthus henteri* based on RAPD markers.

Regarding the genetic variability in *D. nardiformis*, amplification with RAPD markers was very difficult to achieve, only 2 primers as 1225 și OPB-07 allowed amplification with scorable bands (Table 2). By amplification with the primer 1225, only 2 bands were obtained, the first is present only in the individuals from P2 and P3 populations and the second one is present in all individuals independent on the population. Thus, this primer detected low genetic polymorphism between populations, but not within population. By amplification with OPB-07 primers, the RAPD patterns of all individuals are the same, except several individuals from the population P1. The genetic polymorphism detected only with the primer 1225, did not ensure the different clustering of individuals.

In *D. pratensis* ssp. *racovitzae* RAPD amplification was also very difficult, successfully amplification was achieved only with 3 primers as OPE-04, OPB-07 and OPA-13 (Table 2). The amplification with OPE-04 primer generated 3 different fragments, each of them identified in one individual. Amplification with the primer OPB-07 generated 5 fragments that show several differences between individuals. The primer OPA-13 generated only 1 fragment that is present in all individuals. Despite of other species this primer showed no polymorphism in this species. Unfortunately the number of individuals collected form each population is very small and no pertinent conclusion could be showed by these results. The individuals of this species are clustered in 2 groups, in wich are mostly the individuals from population P1. The individual from population P2 belongs to the first cluster (Fig. 6).




Figure 6. Dendrogram illustrating similarities between the individuals of *Dianthus pratensis* ssp. *racovitzae* based on RAPD markers.

In *D. spiculifolius* the RAPD amplification was achieved with the most of the primers, as 1225, OPF-04, OPE-04, OPB-07, OPA-13, 4A26 and 4A27 (Table 2). By amplification with 1225 primer we obtained 5 fragments that showed high genetic polymorphism of the individuals from both populations. Similar results were obtained with the primer OPF-04, 5 polymorphic fragments being identified. With the primer OPE-04, also 5 fragments were obtained, this primer showed genetic polymorphism within populations. By amplification with the primers 4A26 and 4A27 low genetic polymorphis was detected, thus the primer 4A26 generated 2 fragments, and the primer 4A27 generated 3 fragments. These primers showed genetic polymorphism between populations. The primer OPB-07 generated 6 fragment that showed genetic polymorphism between and within populations. The highest polymorphism was detected with the primer OPA13, 7 fragments being identified that showed genetic polymorphism within populations. Dendrogram of similarity shows that most of the individuals are clustered together in one group, and other group is formed only by the individuals 1.1, 2.1 and 2.3 (Fig. 7).

Amplification with RAPD primers of *Dianthus tenuifolius* was successfully obtained with the primers 1225, OPE-04, OPF-04, OPB-07 and OPA-13 (Table 2). The primer 1225 generated 2 fragment, the first fragment is present in all individuals and the second is present only in the individuals from Rodnei Mt. By amplification with OPE-04 primer, 4 fragments were obtained having a polymorphic distribution in the individuals belonging to different populations. The genetic polymorphism of the population P2 is very low. The amplification with the primer OPF-04 generated 5 fragments. The amplification with the primer OPB-07 showed only 2 fragments, the first being present in all individuals and the last one only in the individual 2.3.




Figure 7. Dendrogram illustrating similarities between the individuals of *Dianthus spiculifolius* based on RAPD markers.

The primer OPA-13 generated 4 fragments with polymorphic distribution between individuals. The dendrogram of similarities showed 2 groups, one of them containing the individuals 1.3 and 2.1 and the second group containing the individuals 1.1. and 2.2. The other individuals are more different from those included in these clusters (Fig. 8).




Figure 8. Dendrogram illustrating similarities between the individuals of *Dianthus tenuifolius* based on RAPD markers.

Despite of many morphological (Farsi *et al.* 2013) and cytological (Jafari and Behroozian, 2010) studies on different species of *Dianthus*, there are only limited information on the genetic diversity in the wild populations of these species. There are only few studies regarding the genetic diversity of *D. caryophyllus* using RAPD markers (Su Yeons, 2002) or *D. polylepis* and *D. crinitus* (Behroozian *et al.*, 2013). Recent works revealed low genetic variability of 3 endemic and endangered species of *Dianthus* from Romania as *D. spiculifolius* Schur; *D. giganteus* d'Urv. subsp. *banaticus* (Heuff) and *D. callizonus* (Cristea *et al.* 2014; Jarda *et al.* 2014; Gabel *et al.* 2016).

Our preliminary data obtained with RAPD markers showed low level of polymorphism in the populations of the 8 *Dianthus* species. RAPD markers usually revealed high polymorphism even in the situation when other situs specific markers as SSR showed low polymorphism (Safari *et al.*, 2013; Jarda *et al.*, 2014) but was not the same in our study. Our results are in concordance with other data showing low diversity and polymorphism of *Dianthus* species from Iberian Peninsula (Balao *et al.*, 2010) and Iran (Farsi *et al.*, 2013) and to other data revealing low genetic polymorphism of plants having small populations isolated in fragmented mountain habitats (Duminil *et al.*, 2007). Mountain and alpine plants have to cope with harsh environmental conditions and usually the low genetic diversity is compensated by clonal propagation, which produces rapid, but spatially limited spread of genotypes (Young *et al.*, 2002; Gabel *et al.*, 2016). This is also the situation of *Dianthus* species grown in severe environmental conditions (Stöcklin, 1992).

Conclusions

Genetic diversity of *Dianthus* species revealed by RAPD markers is generally low. Out of 9 primers used in this study, only 6 generated reproducible and polymorphic patterns, useful for preliminary analysis of genetic polymorphism within and between populations.

Acknowledgements. This work was supported by the projects 31-008/2007 and 71/2014, Partnerships in priority areas - PN II, financed by Romanian Ministry of National Education (CNMP and UEFISCDI). Many thanks to Dr. Mihai Pușcaș Dr. Adrian Oprea, Dr. Gheorghe Coldea, Dr. Tudor Ursu and Dr. Liviu Filipaș for collecting the plant material.

REFERENCES

- Balao, F., Valente, L. M., Vargas, P., Herrera, J., Talavera, S. (2010) Radiative evolution of polyploid races of the Iberian carnation *Dianthus broteri* (Caryophyllaceae), *New Phytol.* **187**:542-551

- Behroozian, M., Jafari, A., Farsi, M. (2013) RAPD analysis of genetic variation within and among natural population of two species of *Dianthus* L. (Caryophyllaceae) in NE Iran, *Iran. J. Bot.* **19** (2):194-201
- Breinholt, J. W., Buren, V. R., Kopp, O. R., Stephen, C. L. (2007) Population genetic structure of an endangered Utah endemic, *Astragalus ampullarioides* (Fabaceae), *Am. J. Bot.* **96**: 661-667
- Butiuc-Keul, A. L., Şuteu, A., Munteanu-Deliu, C., Deliu, C. (2001) Study on the in vitro preservation of *Dianthus spiculifolius*, Schur., *Contrib. Bot.* **36**: 137-147
- Butiuc-Keul, A. L. (2006) *Markeri moleculari utilizați în genetică și biotehnologia vegetală* [in Romanian], Mega, Cluj-Napoca, pp. 17-57
- Butiuc-Keul, A. L. (2014) *Biotehnologie generală* [in Romanian], Presa Univeritară Clujeană, Cluj-Napoca, pp. 253-267
- Butiuc-Keul, A. L., Farkas, A., Cristea, V. (2016) Genetic stability assessment of in vitro plants by molecular markers, *Studia UBB Biologia* **LXI**(1):107-114
- Ciocărlan, V. (2009) *Pictured Flora of Romania* [in Romanian], Ceres, Bucureşti.
- Cristea, V., Puşcaş, M., Miclăuş, M., Deliu, C. (2006) Conservative micropropagation of some endemic or rare species from the *Dianthus* genus, *Acta Hort.* **725**: 357-364
- Cristea, V., Deliu, C., Oltean, B., Butiuc-Keul, A. L., Brummer, A., Albu, C., Radu, G.L. (2009), Soiless culture for pharmaceutical use and biodiversity conservation, *Acta Hort.* **843**:157-164
- Cristea, V., Palada, M., Jarda, L., Butiuc-Keul, A. L. (2013) Ex situ in vitro conservation of *Dianthus spiculifolius*, endangered and endemic plant species, *Studia UBB Biologia*, **58**(1):57-69
- Cristea, V., Brummer, A. T., Jarda, L., Miclăuş, M. (2010) In vitro culture initiation and phytohormonal influence on *Dianthus henteri*-a Romanian endemic species, *Rom Biotechnol. Lett.* **25**:25-33
- Cristea, V., Crăciunaş, C., Marcu, D., Palada, M., Butiuc-Keul, A. L. (2014) Genetic stability during *in vitro* propagation of *Dianthus spiculifolius* Schur, *Prop. Ornament. Plants* **14**:26-31
- Doyle, J., Doyle, J. L. (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue, *Phytochem. Bull.* **19**:11-15
- Duminil, J., Fineschi, S., Hampe, A., Jordano, P., Salvini, D., Vendramin, G. G., Petit, R. J. (2007) Can population genetic structure be predicted from life-history traits. *Am. Naturalist* **169**:662-672
- Farsi, M., Behroozian, M., Vaezi, J., Joharchi, M.R., Memariani, F. (2013) The evolution of *Dianthus* polylepis complex (Caryophyllaceae) inferred from morphological and nuclear DNA sequence data: one or two species? *Plant Syst. Evol.* **299**:1419-1431
- Gabel, A. R., Sattler, J., Reisch, C. (2016) Genetic variation and performance of the alpine plant species *Dianthus callizonus* differ in two elevational zones of the Carpathians, *Alp. Bot.* doi:10.1007/s00035-016-0177-3.
- Halmagyi, A., Butiuc-Keul, A. L. (2007) *Conservarea resurselor genetice vegetale* [in Romanian], Todesco, Cluj-Napoca, pp. 122-129
- Jarda, L., Butiuc-Keul, A. L., Höhn, M., Pedryć, A., Cristea, V. (2014) Ex situ conservation of *Dianthus giganteus* d'Urv. subsp. *banaticus* (Heuff.) Tutin by in vitro culture and assessment of somaclonal variability by molecular markers, *Turk. J. Biol.* **38**:21-30

- Johnson, T., Cruse-Sanders, J. M., Pullman, G. S. (2012) Micropropagation and seed cryopreservation of the critically endangered species Tennessee yellow-eye grass, *Xyris tennesseensis* Kral, *In Vitro Cell Dev. Biol. Plant* **48**:369-376
- Luan, S. S., Chiang, T. S., Gong, X. (2006) High genetic diversity vs. low genetic differentiation in *Nouelia insignis* (Asteraceae), a narrowly distributed and endemic species in China, revealed by ISSR fingerprinting, *Ann. Bot.* **98**:583-589
- Marriott, M., Sarasan, V. (2010) Novel micropropagation and weaning methods for the integrated conservation of a critically endangered tree species, *Medusagyne oppositifolia*, *In Vitro Cell Dev. Biol. Plant* **46**:516-523
- Primack, R. B., Pătroescu, M., Rozyłowicz, L., Iojă, C. (2008) *Fundamentele conservării diversității biologice*, AGIR, București, pp. 343-375
- Safari, S., Mehrabi, A. A., Safari, Z. (2013) Efficiency of RAPD and ISSR markers in assessment of genetic diversity in *Brassica napus* genotypes, *Int. J. Agri. Crop Sci.* **5**:273-279
- Stöcklin, J. (1992) Differences in the location of subcotyledonary buds among *Epilobium angustifolium* L., *E. dodonaei* Vill and *E. fleischeri* Hochst (Onagraceae) and effects on architecture and population structure, *Bot. J. Linn. Soc.* **108**:35-47
- Su Yeons, K. (2002) Genetic relationship among Korean *Dianthus* species based on morphological characteristics and RAPD analysis, http://www.niha.go.kr/report_board/report_board_view.asp?size_tag=&sno=1726&gubun=&p_type=11
- Valente, L. M., Savolainen, V., Vargas, P. (2010) Unparalleled rates of species diversification in Europe, *Proc. Royal Soc London B: Biological Sciences* **277**: 1489-1496
- Young, A. G., Hill, J. H., Murray, B. G., Peakall, R. (2002) Breeding system, genetic diversity and clonal structure in the sub-alpine for *Rutidosia leiolepis* F. Muell. (Asteraceae), *Biol. Conservation* **106**:71-78

***Protracheoniscus vasileradui* – n. sp. (Crustacea, Isopoda, Crinochaeta) in the Romanian fauna**

Nicolae Tomescu^{1,✉} and Lucian Alexandru Teodor¹

SUMMARY. We describe the terrestrial isopod species *Protracheoniscus vasileradui* n. sp. collected from the deciduous forest in the Liuborajdea Valley, Iron Gates (Porțile de Fier) area. We named it *Protracheoniscus vasileradui* n.sp., in the memory of Prof. V. Gh. Radu, who was a renowned specialist in terrestrial isopods.

Keywords: isopod, specific characters, *vasileradui*

Introduction

Between 1967 and 1968 V. Gh. Radu and N. Tomescu conducted studies in the habitats of the planned reservoir and neighbouring areas of the Iron Gates (Porțile de Fier). Over 500 terrestrial isopod individuals were collected in this period (Radu and Tomescu, 1975, p. 45-46). In these samples four terrestrial isopod species were found to be new for science. Three of these were described by Radu and published in Fauna of the SRR, Crustacea, Isopoda, vol. IV, fascicle 13, 1983. One of the species mentioned in the Fauna of Porțile de Fier (1975), belonging to the genus *Protracheoniscus*, could not be described at the time by Radu, due to health issues. We recently found 3 ♂ and 8 ♀ *Protracheoniscus* in the samples collected in that period and we named them *Protracheoniscus vasileradui* n.sp., in the memory of Prof. V. Gh. Radu, who was a renowned specialist in terrestrial isopods.

¹ Babeș-Bolyai University, Faculty of Biology and Geology, Department of Taxonomy and Ecology, Cluj-Napoca, Romania.

✉ **Corresponding author:** Nicolae Tomescu, Babeș-Bolyai University of Cluj-Napoca, Department of Taxonomy and Ecology, 5-7 Clinicilor Str., 400006, Cluj-Napoca, Romania.
E-mail: nicolaetomescu36@gmail.com

Materials and methods

The specimens of *Protracheoniscus vasileradui* were collected with the help of a leaf litter sieve, from leaf litter collected in the deciduous forests of the Liuborajdea Valley. The study sites in the Liuborajdea Valley are located between the localities Moldova Veche and Orşova, between the kilometre stones 83 and 82.

Three males and 8 females of this species were collected. Two males have been dissected and microscopic slides of the taxonomically relevant organs were prepared with Euparal and Canada balsam. The extremity of the endopodites of the first male pleopods have been photographed from the slides in ventral position and laterally in alcohol, before dissection. The males and two females have been photographed dorsally, in alcohol, in order to have a whole-body image and the dorsal colouration. Two females were included in Canada Balsam, the six other females and a male are kept in 70% alcohol. These, the whole biological material and the microscopic slides from N. Tomescu's terrestrial isopod collection are going to be donated to the Zoology Museum, Faculty of Biology and Geology of the Babeş-Bolyai University.

In describing the species *Protracheoniscus vasileradui* we will compare some characteristics with those of the species *Protracheoniscus politus*, a common, widespread species from the Romanian fauna. N. Tomescu (1972) described the ontogenetic postembryonic development of *P. politus*, and Tomescu *et al.* (2016) described the change in the morphology of the endopodites of the first male pleopods in relationship with post-reproductive moulting.

Results

Species description:

Size: males: 5 x 2 – 5.2 x 2 mm, females: 5.2 x 2 – 6 x 2.5 mm (Fig. 1 a, b).

Colour: dark-brown with yellow-orange spots.

Somatic features: cephalon: cephalic lobes poorly developed, large yellowish-orange spots on the cephalon (Fig. 1 c). Pleotelson shorter comparatively to that of the species *P. politus* (Fig. 1 d).

Appendages

Antennae. The second-last segment of the antenna is shorter than the last (Fig. 2 a).

Pereiopods. The seventh pereiopod pair of the male has a straight ventral side of the ischiopodite (Fig. 2 b). Males of the species *P. politus* have a slight curvature of the ventral side of the ischiopodite (Fig. 2 c).

Pleopods. The first pair of the male pleopod exopodites are approximately triangular in shape, with the external side slightly concave (Fig. 3 a) and differing from the shape of the first pair of the male pleopod exopodites in *P. politus* (Fig. 3 e).

Endopodites of the male *P. vasileradui* have a wide basal half, with the external sides slightly curved (Fig. 3 b). The extremities of the endopodites are pointed (Fig. 3 c), with tips that are ventrally curved (Fig. 3 d). In males of the species *P. politus*, the first pair of the pleopod endopodites have a chitinous lobe at their extremities (Fig. 3 f, f'), which is shed with the exuvia during post-reproductive moulting and is regrown during the next year (Tomescu, 1972; Tomescu *et al.*, 2016).

Derivatio nominis: we named the species *Protracheoniscus vasileradui* n.sp., in the memory of Prof. V. Gh. Radu, who was a renowned specialist in terrestrial isopods.



Figure 1. *Protracheoniscus vasileradui* nov. spec. Holotyp, male and female dorsal view: **a.** ♂ 5 x 2 mm, **b.** ♀ 5.5 x 2.2 mm, **c.** cephalic lobes, **d.** pleotelson.




Figure 2. *Protracheoniscus vasileradui* nov. spec. Holotyp, ♂ 5 x 2 mm, **a.** antenna, **b.** pereiopods 7. *Protracheoniscus politus*, ♂ 6.5 x 2.8, **c.** pereiopods 7.




Figure 3. *Protracheoniscus vasileradui* nov. spec. Holotyp, ♂ 5 x 2 mm, **a.** exopod pleopods 1, **b.** endopod pleopods 1, **c.** apex of the endopod pleopods 1 – ventral view, **d.** apex of the endopod pleopods 1 – profile view. *Protracheoniscus politus*, ♂ 6.5 x 2.8, **e.** exopod of pleopods 1, **f.** endopod of pleopods 1, **g.** apex of the endopod pleopods 1.

Discussion

The species *Protracheoniscus vasileradui* n. sp. has special morphological characteristics in comparison to *Protracheoniscus politus*, a wide-spread species in Romania. The habitat, where the specimens of *P. vasileradui* were collected from, deciduous forests of the Liuborajdea Valley, in the Iron Gates (Porțile de Fier) area, seems to be a geomorphologically isolated area, that favoured, in time, the isolation of *P. vasileradui* populations. Research conducted in 1967 and 1968 aimed at recording plant and animal species in the area where a reservoir was planned to be formed, by building the Iron Gates dam on the Danube. The research was extended into the habitats of the valley slopes that were not covered by the water of the reservoir. The forest in the Liuborajdea Valley, where we collected the *P. vasileradui* specimens, is located on such a slope, and we believe these populations have survived until the present day. It is possible that the species is present also on areas in the vicinity of the Liuborajdea Valley, with similar ecological conditions.

Conclusions

The specific morphological differences between *Protracheoniscus vasileradui* n. sp. and *Protracheoniscus politus* C. L. Koch 1841 lie in the shape of the male's first pair of exopodites and endopodites, and that of the male's ischiopodites of the seventh pair of pereopods. These are essential morphological characteristics in the separation of species belonging to the genus *Protracheoniscus*.

REFERENCES

- Radu, G. V., Tomescu, N. (1975) *Fauna de la Porțile de Fier, Isopoda* [in Romanian], Ed. Acad. R. S. R. [in Romanian]
- Radu, G. V. (1983) *Crustacea, Isopoda, Oniscoidea, Oniscidae inferioare* [in Romanian], Fauna R.S.R., **IV**(13), pp. 168
- Radu, G. V. (1985) *Crustacea, Isopoda, Oniscoidea, Crinocheta* [in Romanian], Fauna R.S.R., **IV**(14), pp. 155

- Tomescu, N. (1972) Reproduction and ontogenetic development of *Protracheoniscus politus*, *Rev. Roum. Biol., Zool.*, **17** (1), 31-39
- Tomescu, N., Teodor, L. A., Ferenți, S., Covaciu-Marcov, S. D. (2016) Two *Protracheoniscus* Species (Crustacea, Isopoda, Oniscidea) in Romanian Fauna: Morphology, Ecology and Distribution, *Studia UBB Biologia*, **61**(1): 147-166

Clutch size and egg repeatability in three elusive bird species: Little Bittern (*Ixobrychus minutus*), Little Crake (*Zapornia parva*) and Water Rail (*Rallus aquaticus*) from north-west Romanian populations

Alin David¹, Alexandru Nicolae Stermin^{1,✉} and Eliana Sevianu²

SUMMARY. Data were collected from Little Bittern, Water Rail and Little Crake nests located in North-Western Romania, between April and July of 2002 - 2006 and 2010 - 2012. The repeatability was calculated as intra-class correlation of length, breadth and egg volume coefficients. The total number of active nests considered for the present study was as follows: 43 for Little Bittern, 25 for Water Rail and 9 for Little Crake. The clutch size in Little Bittern ranged from 4 to 7 eggs, in Water Rail from 4 to 10 and in Little Crake from 4 to 8 eggs. The repeatability values of egg parameters varied between 0.844 to 0.860 in Little Bittern, 0.262 - 0.374 in Water Rail and 0.310 - 0.574 in Little Crake. Short-term environmental impact was strong in case of Water Rail and Little Crake, while the genetic component had little influence. For these two species, the low repeatability could be explained by larger numbers of eggs in a clutch, reflecting in turn a longer laying period.

Keywords: breadth, egg volume, laying period, length.

Introduction

Clutch size and egg characteristics vary among and within bird species (Lack, 1947; Klomp 1970; Figuerola and Green, 2005). These variations are important in revealing the individual's fitness (Rockwell *et al.*, 1987, Brisk and Sealy, 1989; Horak *et al.*, 1997) and aspects about species evolution (Martin, 1995; Figuerola and Green, 2005).

¹ Babeș-Bolyai University, Department of Taxonomy and Ecology, Cluj-Napoca, Romania.

² Babeș-Bolyai University, Department of Environmental Science, Cluj-Napoca, Romania.

✉ **Corresponding author:** Alexandru Nicolae Stermin, Department of Taxonomy and Ecology, Babeș-Bolyai University, Cluj-Napoca, Romania,
E-mail: sandu.stermin@yahoo.com

In bird species, clutch size may depend on many factors, such as the effective coverage capacity of the eggs during incubation (Rice and Kenyon, 1962) or the pressure from predators during nesting (in the sense that, the more eggs are being laid, the more exposed to predators they are) (Slagsvold, 1982; Arnold *et al.*, 1987). The chances in offspring survival of large clutches usually decrease, in the sense that as the number of the offspring is larger, the possibility of being detected by predators increases (Cooch, 1961; Hilden, 1964; Lessells, 1986). Other factors influencing the clutch size are: food resources available during egg laying (Lark, 1974) and temperature during egg production (Great Tits, *Parus major* roosting in cooled nestboxes laid eggs 14% smaller than those roosting in heated nestboxes (Nager and Noordwijk, 1992)). In addition, significant effects on clutch size and egg characteristics are reported from parameters like: female age and experience (a statistically significant increase in egg size with age or experience was found in almost half (17/37) of the studies (Christians, 2002)); female mass and size; and other aspects of the phenotype (Christians, 2002).

The egg size and their intraclutch and interclutches variations are important in determining the offspring body size and weight, their survival probability and therefore the reproductive success (Schifferli, 1973; Williams, 1994; Surmacki, 2003). Variations in egg size are the consequences of genetic and environmental action (Surmacki, 2003), repeatability dimensions characterizing environmental conditions and food resources during laying period (Stermin *et al.*, 2008).

By calculating the repeatability of egg size we can split the phenotypic variation in egg size into within-individual and among-individual components. Variation in egg size among individuals is caused by a combination of genetic and environmental differences, while the variation in egg size within an individual during several years is caused by temporary environmental differences between reproductive attempts (Falconer, 1989; Flint *et al.*, 2001).

In this context, analyzing the repeatability in egg size can reveal information about the ecology and biology of bird species. These data bring valuable information about several bird species which are poorly studied, due to their elusiveness and rather inaccessible habitats. Some of the less studied bird species in Europe are Little Bittern *Ixobrychus minutus*, Little Crake *Zapornia parva* and Water Rail *Rallus aquaticus* (Marion *et al.*, 2000; Stermin *et al.*, 2011; Stermin *et al.*, 2013).

Those three species coexist in dense wetland habitats and during their evolution each has adapted to occupy specific microhabitats inside the wetlands (Taylor, 1998; David, 2008; Stermin *et al.*, 2012). Little Bittern and Little Crake are protected species and, along with the Water Rail, are now being affected by habitat loss and fragmentation (Marion *et al.*, 2000; Brambilla *et al.*, 2012). New information about their biology and ecology is important not only in solving the mystery of their elusive life, but also in designing and implementing conservation programs.

Thus, the main goals of our study are: (i) to describe the variation of clutch size and egg parameters in Little Bittern, Water Rail and Little Crake populations from the Fizeş Basin (N-W Romania) and (ii) to calculate the repeatability and compare the results to other studies on other bird species by analyzing them in relation to the effect of the environment (sensus Bańbura and Zieliński, 1998).

Materials and methods

The study area is located in the central part of the Transylvanian Plain, in Romania (24°10' E; 46°50'N). The wetlands surveyed here represent 1/3 from all Transylvanian wetlands. Most wetlands from the Fizeş Basin are man-made and used as fishing ponds; only two of them are natural, and were recently designated "Natura 2000" site (Special Protected Area) (David, 2008). The wetland areas range from 19 to 253 ha, being partially or totally covered by marsh vegetation.

Data were collected between April and July of 2002 - 2006 and 2010 - 2012. During this time, the areas suitable for Little Bittern, Little Crake and Water Rail nesting were checked at intervals of 2 to 5 days. Transect method was used in nest searching (Bibby *et al.*, 2000). The Little Bittern nests were located in Lakes Ştiucii, Borzaş, Năsal, Ţaga Mare, Sucutard I, Roşieni, Tău Popii, Cătina and Legii Pond; the Little Crake nests were found in Lakes Sucutard I and Cătina, while the Water Rail nests were situated on Sic Reedbeds and Lake Ştiucii.

The length (L) and breadth (B) of eggs were measured to the nearest 0.1 mm with digital caliper. The volume was calculated using the following formula: $V = k \times L \times B^2 / 1000$ (where $k = 0.51$) (Hoyt, 1979). The egg weight was measured in the field using a digital scale.

One-way analysis of variance (ANOVA) was carried out in order to obtain variance components. The repeatability was calculated as intra-class correlation coefficients (Lessells and Boag, 1987; Falconer and Mackay, 1995; Sokal and Rohlf, 1995), standard errors for repeatability values were calculated as described in Becker (1992).

Results and discussion

A total of 223 eggs from 43 active Little Bittern nests, 209 eggs from 25 Water Rail nests and 56 eggs from 9 Little Crake nests were measured. The clutch size in Little Bittern ranged from 4 to 7 eggs, in Water Rail from 4 to 10 and in Little Crake from 4 to 8 (Table 1).

Table 1.

Clutch size characteristics in Little Bittern, Water Rail and Little Crane nests located in the Fizeş Basin (SD - standard deviation)

	Little Bittern	Water Rail	Little Crane
Total number of clutches	43	25	9
Minimum number of eggs/clutch	4	4	4
Maximum number of eggs/clutch	7	10	8
Mean of eggs/clutch \pm SD	5.32 \pm 0.74	8.36 \pm 1.55	6.22 \pm 1.39

Egg measurements for Little Bittern, Water Rail and Little Crane are presented in table 2, while repeatability values are depicted in table 3.

Table 2.

Egg measurements from Little Bittern, Water Rail and Little Crane clutches located in the Fizeş Basin (SD - standard deviation)

		Little Bittern (n = 223)	Water Rail (n= 209)	Little Crane (n=39)
Length (mm)	Minimum	30.80	26.59	28.87
	Maximum	37.80	39.59	34.09
	Mean \pm SD	34.92 \pm 1.45	35.92 \pm 1.64	31.01 \pm 1.38
Breadth (mm)	Minimum	22.10	23.88	20.58
	Maximum	29.10	36.93	22.81
	Mean \pm SD	25.55 \pm 0.92	25.96 \pm 1.24	21.76 \pm 0.57
Volume (cm ³)	Minimum	8.81	8.63	6.33
	Maximum	16.15	24.14	8.42
	Mean \pm SD	11.66 \pm 1.14	12.38 \pm 1.42	7.49 \pm 0.58
Weight (g)	Minimum	-	9.9	6.60
	Maximum	-	15.10	8.90
	Mean \pm SD	-	12.43 \pm 0.95	7.71 \pm 0.61

Table 3.

Repeatability (R) of egg length, breadth and volume index in Little Bittern, Water Rail and Little Crake clutches located in the Fizeş Bazin (SE - standard error; Df - degrees of freedom; F - variance ratio; P - significance level)

		R	SE	df	F	P
Little Bittern	Length	0.844	3.055	42	23.374	< 0.001
	Breadth	0.860	1.639	42	6.363	< 0.001
	Volume	0.851	2.225	42	10.932	< 0.001
Water Rail	Length	0.374	9.326	23	8.537	< 0.001
	Breadth	0.355	7.270	23	9.863	< 0.001
	Volume	0.262	15.824	23	28.212	< 0.001
Little Crake	Length	0.574	10.771	8	16.092	< 0.001
	Breadth	0.517	3.288	8	7.294	< 0.001
	Volume	0.310	3.917	8	24.83	< 0.001

Significant statistical differences between species were found regarding egg measurements: length ($F(2.468) = 196.60$, $df = 2$, $P < 0.001$), breadth ($F(2.468) = 262.95$, $df = 2$, $P < 0.001$) and volume ($F(2.468) = 252.80$, $df = 2$, $P < 0.001$). Larger eggs were recorded in Water Rail, while Little Bittern and Little Crake had smaller eggs.

Previous studies reported that the Water Rail clutch size ranged from 5 to 16 eggs, with an average of 6 - 11 eggs (Flegg and Glue, 1973; Taylor, 1998). In the present study, the smallest clutch contained 4 eggs and the largest one 10 eggs, most of the clutches having 8 - 9 eggs. Moreover, the average egg size corresponded to those described by other authors, who observed egg sizes of $32 - 40 \times 24.1 - 27.2$ mm (Cramp and Simmons, 1980; Taylor, 1998). From this point of view, no differences were depicted between European populations and ones from the Fizeş Basin. A similar situation was recorded for Little Crake: its clutch size was reported to range from 4 to 11 eggs (Cramp and Simmons, 1980; Taylor, 1998), with an average size of 7 - 9 eggs. In this study, the 4 egg clutch was a replacement of the first clutch, destroyed by flooding. The average egg size was quite similar to those described in other studies, ranging from $28.5 - 32 \times 12.2 - 22.2$ mm (Dombrowski, 1912) or from $27.5 - 33.5 \times 19 - 23$ mm, with a weight of 6.3 - 8.7 g (Taylor, 1998).

In previous studies on Little Bittern, the clutch size ranged from 4 to 7 eggs (Samraoui *et al.*, 2012); 1 to 6 eggs (Pardo-Cervera *et al.*, 2010), 4 to 9 eggs (Cramp and Simmons, 1977), or 3 to 7 eggs (Martínez-Abraín, 1994). This clutch

size variation could be a consequence of various habitat quality, since Little Bittern was described as an area-sensitive species (Pezzo and Benocci, 2001; Benassi *et al.*, 2009). The variation of egg sizes in the present study was comparable to previous literature, where the means for egg length were 35.1 ± 1.3 mm; for egg breadth 25.7 ± 0.7 mm, and for egg volume 11.8 ± 0.9 cm³ (Samraoui *et al.*, 2012), and the mean egg size 34.2 mm \times 25.6 mm (range $30.9 - 37.3 \times 23.6 - 26.9$ mm) (Pardo-Cervera *et al.*, 2010).

Regarding the repeatability, the lowest values were recorded for volume and weight. These low values are normal in volume analysis, since it is a parameter that increases approximately with the cube of the linear size. On the other hand, weight is a parameter proportional to volume and density, and is therefore less related to linear dimensions, thus having a different effect on the repeatability calculation.

Repeatability values in Little Crake (0.574 - 0.260) and Water Rail (0.374 - 0.262) were low compared to those in Little Bittern (0.860 - 0.844), and those calculated in other birds (Boag and van Noordwijk, 1987; Hendricks, 1991; Potti, 1993; Bańbura and Zieliński, 1998; Jerzak *et al.*, 2000; Tryjanowski *et al.*, 2001; Christians, 2002; Zduniak and Antczak, 2003), which are generally higher than 0.6 (Christians, 2002). However, lower repeatability values were reported in the literature, in Greater Scaup, *Aythya marila* and American Oystercatcher, *Haematopus palliatus* (0.36 in both cases) (Nol *et al.*, 1984; Flint and Gran, 1999). Higher repeatability values were found in Redshank *Tringa tetanus* (Thompson and Hale, 1991), Northern Pintail *Anas acuta* (Flint and Grand, 1996) and Canada Goose *Branta canadensis* (Leblanc, 1989): 0.87, 0.89 and 0.92, respectively.

These findings highlighted the fact that in Water Rail and Little Crake the environmental impact was strong on short term and the genetic component had very little influence.

The low repeatability in Water Rail and Little Crake could be an effect of the large clutch size, which means a long laying period, compared with Little Bittern (with smaller clutch size and a shorter laying period). During a longer laying period, birds may face environmental variations that affect accessibility to food and thus the amount of resources and nutrients invested into an egg. Previous studies inferred that in some bird species habitat quality and food supply may affect the eggs (Christians, 2002). On the other hand, other results show that food availability may have no effect on egg size (Arnold *et al.*, 1991; Bolton *et al.* 1993; Jager *et al.*, 2000).

A higher repeatability in Little Bittern could be caused by smaller clutch size, that implied a short laying period of time, and a lower probability of environmental changes.

The wetland habitats have rapidly changing parameters in terms of the water level, vegetation density and availability of food resources (Dyrcz and Zdunek, 1996). In wetlands, adaptability is a condition of survival, in contrast to birds living

in urban environments, which have a higher rate of repeatability, living under more constant environmental conditions and where food resources are constant over longer periods (Surmacky *et al.*, 2003).

REFERENCES

- Arnold, T. W., Alisauskasand, R. T., Ankney, C. D. (1991) Egg composition of American coots in relation to habitat, year, laying date, clutch size, and supplemental feeding. *Auk* **108**: 532-547
- Bañbura, J., Zieliński, P. (1998) An analysis of egg-size repeatability in Barn Swallows *Hirundo rustica*. *Ardeola* **45**: 183-192
- Benassi, G., Battisti, C., Luiselli, L., Boitani, L. (2009) Areasensitivity of three reed bed bird species breeding in Mediterranean marshland fragments, *Wetlands Ecol Manage* **17**: 555-564
- Bibby, C. J., Burgess, N. D., Mustoe, S. H. (2000) *Bird census techniques*, London: Academic Press
- Boag, P. T., van Noordwijk, A. J. (1987) Quantitative genetics. In: Buckley, P. A., Cooke, F. (ed.) *Avian genetics*, Academic Press, London, pp. 45-78
- Bolton, M., Monaghan, P., Houston, D. C. (1993) Proximate determination of clutch size in lesser black-backed gulls: the roles of food supply and body condition, *Canadian Journal of Zoology* **71**: 273-279
- Christians, J. K. (2002) Avian egg size: variation within species and inflexibility within individuals, *Biological Reviews* **77**: 1-26
- Ciach, M. (2011) Habitat-related differences in egg size in the Spur-Winged Lapwing *Vanellus spinosus*. *Ardeola* **58**: 335-341
- David, A. (2008) *Ecologia populațiilor de păsări din Câmpia Fizeșului*. Presa Universitară Clujeană, Cluj-Napoca [in Romanian]
- Dombrowski, R. R. (1912) *Ornis Romaniae: die Vogelwelt Rumänien's systematisch und biologisch-geographisch beschrieben*, Staatsdruckerei, Bukarest [in German]
- Dyrcz, A., Zdunek, W. (1996) Potential food resources and nestling food in the Great Reed Warbler *Acrocephalus arundinaceus* and Reed Warbler *Acrocephalus scirpaceus* at Milicz fish-ponds, *Ptaki Śląska* **11**: 123-132
- Falcone, D. S. (1989) *Introduction to Quantitative Genetics*, John Wiley and Sons, New York
- Figuerola, J., Green, A. (2005) A comparative study of egg mass and clutch size in the Anseriformes. *J Ornithol*. DOI 10.1007/s10336-005-0017-5
- Fint, P. L., Grand, J. B. (1996) Variation in egg size of the Northern Pintail, *Condor* **98**: 162-165
- Flegg, J. J. M., Glue, D. E. (1973) A Water Rail study, *Bird Study* **20**: 69-80
- Flint, P. L., Grand, J. B. (1999) Patterns of variation in size and composition of Greater Scaup eggs: are they related? *Wilson Bulletin* **111**: 465-471
- Flint, P. L., Rockwell, R. F., Sedinger, J. S. (2001) Estimating repeatability of egg size, *Auk* **118**: 500-503

- Hendricks, P. (1991) Repeatability of size and shape of American Pipit eggs, *Can. J. Zool.* **69**: 2624-2628
- Hoyt, D. E. (1979) Practical methods for estimating volume and fresh weight of birds eggs, *Auk* **96**: 73-77
- Jager, T. D., Hulscher, J. B., Kersten, M. (2000) Egg size, egg composition and reproductive success in the oystercatcher *Haematopus ostralegus*, *Ibis* **142**: 603-613
- Jerzak, L., Bocheński, M., Kuczyński, L., Tryjanowski, P. (2000) Repeatability of size and shape of eggs in the urban Magpie *Pica pica* (Passeriformes: Corvidae) population, *Acta zool Cracov* **43**: 165-169
- Leblanc, Y. (1989) Variation in size and eggs of captive and wild Canada Geese, *Ornis Scandinavica* **20**: 93-98
- Marion, L., Ulenaers, P., Vanvessem, J. (2000) Herons in Europe, In: Kushalan, J. A., Hafner, H. (eds). *Heron Conservation*. Acad. Press, London
- Martinez-Abraín, A. (1994) Notes on the biology of Little Bittern *Ixobrychus m. minutus* during the breeding season in eastern Spain, *Ardeola* **41**: 169-171
- Nager, R. G., Noordwijk, A. J. V. (1992) Energetic limitation in the egg-laying period of Great Tits, *Proceedings of the Royal Society of London Series B* **249**: 259-263
- Nol, E., Baker, A. J., Cadman, M. D. (1984) Clutch initiation dates, clutch size, and egg size of the American Oystercatcher in Virginia, *Auk* **101**: 855-867
- Pardo-Cervera, F., Sørensen, I. H., Jensen, C., Ruiz, X., Sanchez Alonso, C. (2010) Breeding biology of the Little Bittern *Ixobrychus minutus* in the Ebro delta (NE Spain), *Ardeola* **57**: 407-416
- Pezzo, F., Benocci, A. (2001) Spatial behavior of the Little Bittern (*Ixobrychus minutus*): implications for conservation, *Avocetta* **25**: 78
- Potti, J. (1993) Environmental, ontogenetic, and genetic variation in egg size of Pied Flycatchers, *Can. J. Zool.* **71**: 1534-1542
- Samraoui, F., Nedjah, R., Boucheker, A., Alfarhan, A. H., Samraoui, B. (2012) Breeding ecology of the Little Bittern *Ixobrychus minutus* in northeast Algeria, *Bird Study*, iFirst, 1-8, DOI:10.1080/00063657.2012.733335
- Stermin, A. N., Pripon, L. R., David, A., Coroiu, I. (2011) Wetlands management for Little Crake (*Porzana parva*) conservation in a “Natura 2000” site, *International Conference for Environmental Sciences and Development, IPCBEE* **4**: 91-94
- Stermin, A. N., David, A., Sevianu, E. (2013) An Evaluation of Acoustic Monitoring Methods for a Water Rail (*Rallus aquaticus*) Population in a Large Reed Bed, *Waterbirds* **36**: 463-469
- Surmacki, A., Stępiewski, J., Zduniak, P. (2003) Repeatability of egg dimensions within the clutches of Bearded Tit *Panurus biarmicus*, *Acta Ornithologica* **38**: 123-127
- Thompson, P. S., Hale, W. G. (1991) Age-related reproductive variation in the Redshank *Tringa totanus*, *Ornis Scandinavica* **22**: 353-359
- Tryjanowski, P., Kuczyński, L., Antczak, M., Skoracki, M., Hromada, M. (2001) Within-clutch repeatability of egg dimensions in the jackdaw *Corvus monedula*: a study based on a museum collection, *Biologia Bratislava* **56**: 211-215
- Zduniak, P., Antczak, M. (2003) Repeatability and within-clutch variation in egg dimensions in a Hooded Crow *Corvus corone cornix* population, *Biol. Lett.* **40**: 37-42